

Recommendations for the Control of Gastroenteritis Outbreaks in Long-Term Care Homes

**Recommendations for Long-Term Care Homes and
Public Health Unit Staff**

Population and Public Health Division
Ministry of Health and Long-Term Care
March 2018

This document reflects the committee's consensus position on prudent practice and is made available as a resource to public health and health-care providers.

All or part of this document may be reproduced without permission with the following acknowledgement:

Ministry of Health and Long-Term Care/Population and Public Health Division
Toronto, Canada, 2018

© Queen's Printer for Ontario

Ministry of Health and Long-Term Care
Copies of this report can be obtained from
INFOline: 1-866-532-3161

TTY 1-800-387-5559

Table of Contents

Executive summary	6
Acknowledgements	7
Abbreviations	10
Glossary	12
1 Introduction	17
1.1 Purpose	17
1.2 The Role of Public Health	19
1.3 The Role and Responsibilities of the Long-Term Care Homes	20
1.4 Types of Gastroenteritis Outbreaks	20
1.4.1 Outbreaks Caused by Viruses.....	20
1.4.2 Outbreaks Caused by Bacteria and Parasites	21
2 Preparation – Infection Control and Outbreak Policies and Procedures	22
2.1 Outbreak Preparedness	23
2.2 The Infection Prevention and Control Program – Disease Prevention	25
2.2.1 Routine Practices	25
2.3 Additional Precautions	30
2.3.1 Contact Precautions	30
2.3.2 Other Disease Prevention Strategies	30
2.3.3 Education of Staff and Volunteers.....	32
2.3.4 Education of Residents and Visitors.....	33
3 Surveillance and Outbreak Detection	34
3.1 Surveillance	34
3.1.1 Target Groups for Surveillance	34
3.2 Outbreak Detection	35
3.2.1 Infectious Gastroenteritis Case Definition.....	36
3.2.2 Outbreak Definition	36
3.2.3 Suspected Gastroenteritis Outbreak Definition	37
3.2.4 Gastroenteritis Outbreak Definition	37
4 Managing Gastroenteritis Outbreaks	37
4.1 Control Measures for a Suspected Gastroenteritis Outbreak	37

4.2	Steps in Outbreak Management	38
4.2.1	Assess the Outbreak.....	38
4.2.2	Implement General Infection Prevention and Control Measures	38
4.2.3	Consult with the Public Health Unit	38
4.2.4	Declare an Outbreak	39
4.2.5	Notify Appropriate Individuals/Agencies	39
4.2.6	Call an Outbreak Management Team Meeting	40
4.2.7	Monitor the Outbreak on an Ongoing Basis	42
5	Control Measures For Specific Groups	43
5.1	Control Measures for Residents	43
5.1.1	Admissions and Returns from Absences.....	45
5.1.2	Medical and Other Appointments	46
5.1.3	Transfer to Hospital, Other Facility or Urgent Appointment	46
5.2	Control Measures for Staff and Volunteers	46
5.2.1	Well Staff.....	46
5.2.2	ILL Staff.....	47
5.3	Control Measures for Visitors and During Communal Activities	47
5.3.1	Modified Visitor Access	47
5.3.2	Communal Meetings and Other Activities	48
6	Declaring the Outbreak Over and Termination of Outbreak Control Measures	49
6.1	Criteria for Declaring an Outbreak Over	49
6.2	Review the Outbreak	50
6.3	Complete the Outbreak Investigation File	50
7	Investigation and Management of Food-Borne Outbreaks	51
7.1	Recommended Food Handling Policies and Procedures	51
7.1.1	Food-Handling Staff	52
7.1.2	Records of Food Suppliers.....	52
7.1.3	Retaining Food Samples.....	52
7.1.4	Temperature Records of Potentially Hazardous Food	52
7.1.5	Catered Food and Food Brought in by Families.....	53
7.1.6	Common Servery or Kitchenette	53
7.1.7	Feeding Assistance.....	54
7.1.8	Dishwasher Temperature and Sanitizing Records.....	54

7.1.9 Kitchen Equipment Installation and Maintenance.....	54
7.2 Food-borne Illness Investigation.....	54
7.2.1 Questionnaires.....	55
7.2.2 Clinical Samples.....	55
7.2.3 Food Service Operation	55
7.2.4 HACCP Investigation	55
7.2.5 Provision of Alternative Sources of Food and Water.....	56
7.2.6 Summary Report.....	56
References and further information	57
Appendix 1: Instructions for Feces Specimen Collection During Outbreaks	61
Appendix 2: Routine Practices, Additional Precautions, and reporting requirements for selected pathogens or conditions associated with gastroenteritis outbreaks	64
Appendix 3: Routine Practices for all Health Care Settings.....	67
Appendix 4: Signage for entrance to room of a resident requiring Contact Precautions in non-acute care facilities.....	68
Appendix 5: Hand Hygiene Fact Sheet for Health Care Settings	69
Appendix 6: Recommended steps for putting on and taking off personal protective equipment (PPE)	70
Appendix 7: Donning and removal of personal protective equipment (PPE)	72
Appendix 8: Cleaning checklist for an ill resident’s room during an outbreak	73
Appendix 9: Sample screening-policies sign to use during outbreaks.....	75
Appendix 10: Sample surveillance sheet.....	76
Appendix 11: Quick reference guide: suspected outbreak	77
Appendix 12: Steps in the prevention and management of gastroenteritis outbreaks	80
Appendix 13: Gastroenteritis outbreak: sample line-listing form.....	81
Appendix 14: Position Statement: Recommended Length of Exclusion for Cases Associated with Norovirus Outbreaks and When to Declare Norovirus Outbreaks Over.....	82
Appendix 15: Sample Transfer & Return Algorithm for Use During Outbreaks	86
Appendix 16: Epidemiology of selected pathogens associated with gastroenteritis outbreaks	88

Executive summary

Gastroenteritis outbreaks in institutions became a Reportable Disease under *the Health Protection and Promotion Act* in 1991. The increased person-to-person contact and presence of a population with weakened immune systems increases the risk of gastroenteritis outbreaks in Long-Term Care Homes (LTCHs). Gastroenteritis outbreaks continue to occur and to control them; LTCHs require the development of effective infection control programs.

Recommendations for the Control of Gastroenteritis Outbreaks in Long-Term Care Homes (MOHLTC, 2018) replaces Control of Gastroenteritis Outbreaks in Long-Term Care Homes (MOHLTC, 2013). The purpose of the document is to assist LTCHs before, during, and after a gastroenteritis outbreak and to minimize illness, hospitalization, and death, related to gastroenteritis outbreaks in LTCHs.

This document will provide useful and practical information that will help LTCHs:

1. Develop outbreak management policies and procedures and infection control programs.
2. Understand the importance of surveillance, early identification, and isolation of cases to prevent disease spread.
3. Educate their staff on policies and procedures and the importance of Routine Practices.
4. Respond to and manage a gastroenteritis outbreak should one occur.
5. Develop procedures for managing outbreaks associated with food.

The recommendations presented are based on the most current, evidence-based literature, clinical knowledge, trends, and expert consensus on prevention, detection, management, and control of gastroenteritis outbreaks. Included are quick reference appendices that outline procedures for managing specific micro-organisms (pathogens) and summaries of what to do if an outbreak is suspected and what to do during an outbreak.

Changes in the February 2018 version of the Recommendations for the Control of Gastroenteritis Outbreaks in Long-Term Care Homes are limited to title change and reference to the Standards.

Acknowledgements

The Ministry of Health and Long-Term Care would like to acknowledge the contribution and expertise of the working group that developed *Control of Gastroenteritis Outbreaks in Long-Term Care Homes* in 2011 and provided updates to the 2013 version.

Karen Beckermann

Scheduling and Logistics Supervisor East York, Vaccine Preventable Diseases
Toronto Public Health

Risa Cashmore

Infection Control Consultant, Central West Infection Control Network
Public Health Ontario

Joanne Dow

Public Health Nurse, Infectious Disease Control Team
Middlesex-London Health Unit

Lisa Fortuna *

Director, Communicable Disease Prevention and Control
Public Health Ontario

David Fraser

Food and Rabies Coordinator, Health Protection Service
Simcoe-Muskoka District Health Unit

Edwina Gracias

Manager, Health Protection
Peterborough County-City Health Unit

Judy de Grosbois

Public Health Inspector
Perth District Health Unit

Lucie Imbiscuso

Public Health Inspector
Wellington Dufferin Guelph Public Health Unit/Canadian Institute of Public Health Inspectors (Ontario Branch)

Manisa Jiaravuthisan

Associate Director, Communicable Diseases Control
Toronto Public Health

Lois Lacroix

Manager, Infectious Disease Program
Niagara Region Public Health Department

Andre LaFlamme

Manager, Environmental Health
Niagara Region Public Health Department

Joseph Y. Lam

Supervisor, Outbreak Management
Ottawa Public Health

Bill Limerick

Retired, Director of Environmental Health, Director of Health Protection
Northwestern Health Unit

Marina Lombos

Head Technologist, Laboratory Enteric & Molecular Surveillance
Public Health Ontario

Anne Maki

Manager, Enteric, Environmental, Molecular Surveillance and STI
Public Health Ontario

Roman Malanczyj

Senior Public Health Inspector
Durham Region Health Department

Dr. Dean Middleton *

Senior Public Health Epidemiologist, Enteric, Zoonotic and Vector-Borne Diseases
Public Health Ontario

Francine Paquette

Infection Control Consultant, South Western Ontario Infection Control Network
Public Health Ontario

Brenda Smith

Network Coordinator, Central West Infection Control Network
Public Health Ontario

Dave Stronach

Retired, Manager
Halton Region Health Department

Debbie Valickis

Infection Control Specialist, Communicable Disease Division
Peel Public Health

Yvonne Whitfield *

Senior Program Consultant, Enteric, Zoonotic and Vector-Borne Diseases
Public Health Ontario

Anne-Luise Winter

Senior Epidemiologist, Communicable Disease Prevention and Control
Public Health Ontario

* At the time of the development of the original document in 2011 these contributors were affiliated with the Ministry of Health and Long-Term Care, Public Health Division.

Post-Panel Contributors

Jane Carruthers

Long-Term Care Homes Inspector
Ministry of Health and Long-Term Care

Richard Hayden

Long-Term Care Homes Inspector
Ministry of Health and Long-Term Care

Dr. Zsuzsanna Rajda

Senior Program Consultant, Public Health Policy and Programs Branch
Public Health Division, Ministry of Health and Long-Term Care

Dr Erika Bontovics

Manager, Public Health Policy and Programs Branch
Public Health Division, Ministry of Health and Long-Term Care

Stephen Moore

Manager, Enteric, Zoonotic and Vector-borne Diseases
Public Health Ontario

Dr. Doug Sider

Medical Director, Infectious Diseases
Public Health Ontario

Abbreviations

ABHR

Alcohol-Based Hand Rub

CDI

Clostridium difficile Infection

CHICA

Community Health and Infection Control Association

HACCP

Hazard Analysis Critical Control Point

HAI

Health Care-Associated Infections

HPPA

Health Protection and Promotion Act

ICP

Infection Prevention and Control Professional

IPC

Infection Prevention and Control

LHIN

Local Health Integration Network

LTCH

Long-Term Care Home

LTCHA

Long-Term Care Homes Act

MOHLTC

Ministry of Health and Long-Term Care

OHA

Ontario Hospital Association

OHSA

Occupational Health and Safety Act

OMA

Ontario Medical Association

OMT

Outbreak Management Team

PIDAC

Provincial Infectious Diseases Advisory Committee

PHAC

Public Health Agency of Canada

PHI

Public Health Inspector

PHF

Potentially Hazardous Food

PHN

Public Health Nurse

PPE

Personal Protective Equipment

RICN

Regional Infection Control Network

Glossary

Additional Precautions –

These precautions (i.e. Contact Precautions, Droplet Precautions and Airborne Precautions) are carried out in addition to Routine Practices when infections caused by organisms transmitted by these routes are suspected or diagnosed. They include the physical separation of infected or colonized residents from other individuals and the use of barriers (e.g. gowns, gloves, masks) to prevent or limit the transmission of the infectious agent, from colonized or infected individuals, to those who are susceptible to infection or to those who may spread the agent to others.

Alcohol-based hand rub (ABHR) –

A liquid, gel or foam formulation of alcohol (e.g., ethanol, isopropanol) which is used to reduce the number of micro-organisms on hands in situations when the hands are dry and not visibly soiled. ABHRs are less time-consuming to use than washing with soap and water.

Attack rate –

The occurrence of disease observed among a defined population over a limited period of time.

Baseline –

The normal level or presence of a disease or infectious agent within a given geographic area or a population group and time period.

Case –

A person with the particular illness or disease, usually fitting the case definition.

Case definition –

A set of criteria for determining who should be classified as a case. The definition is comprised of clinical information and should include epidemiological information related to time, place, and person.

Cohorting –

Cohorting of residents: Grouping of residents who present either with the same set of symptoms or are asymptomatic. Cohorting of staff: Grouping of staff to care for a specific group of residents or to assign them to a floor/unit that either contains or does not contain active cases.

Common-source outbreak –

A type of outbreak that occurs when individuals are exposed to a point-source of infection at the same time.

Contact Precautions –

A type of Additional Precautions to reduce the risk of transmitting infectious agents via contact with an infectious person. Contact Precautions are used in addition to Routine Practices.

Contact time –

The length of time a surface is exposed to a disinfectant in order for the disinfectant to be effective against micro-organisms.

Contract worker –

Workers from an outside agency. These workers include health-care workers, maintenance and other workers or those who carry on activities in resident-care areas or come into contact with residents, such as hairdressers.

Control measure –

Any action or activity that can be used to prevent or stop an outbreak. Control measures for gastroenteritis outbreaks are primarily focused on reducing additional exposure.

Cross-contamination –

The transfer of pathogens from one food item to another during food preparation through cooking equipment, utensils or the hands of food handlers.

Employees (on payroll) –

This includes all persons who receives a direct paycheck from the reporting facility (i.e., on the facility's payroll), regardless of clinical responsibility or patient contact. This category should not include staff on long-term leave (e.g. maternity, paternity, disability).

Endemic –

The usual presence of a disease or infectious agent within a given geographic area or a population group. Usually expressed as a rate of prevalence.

Fecal-oral transmission –

Transmission of micro-organisms such as bacteria, viruses and parasites from feces into the mouth, through contaminated hands, food, water or objects.

Food handler –

A person who directly handles or prepares food.

Fomite –

An inanimate object that may be contaminated with infectious organisms and serve in their transmission.

Gastroenteritis –

Inflammation of the stomach and intestines that usually causes diarrhea and/or vomiting.

Hazard Analysis Critical Control Point (HACCP) –

A science-based, systematic approach of identifying, evaluating, and controlling food- safety hazards. HACCP is designed to prevent, reduce or eliminate potential biological, chemical and physical food-safety hazards, including those caused by cross- contamination.

Health-care setting –

Any location where health care is provided, including settings where emergency care is provided, hospitals, LTCHs, outpatient clinics, community health centres and clinics, physician offices, dental offices, and home health care.

Incubation period –

The interval from the time an individual is infected to the time when symptoms first appear.

Infection prevention and control committee –

A group that meets regularly to discuss infection-control issues. LTCHs are required to have infection-control committees.

Infection prevention and control professional (ICP) –

A health professional designated to be responsible for infection control programs in the LTCH, in accordance with LTCHA, 2007, S.O. 2007, c. 8 and O. Reg. 79/10. The ICP should possess expertise and additional training in infection prevention and control. In a LTCH, there is a designated staff member to coordinate the program who has education and experience in infection prevention and control practices (including infectious diseases, cleaning and disinfection, data collection and trend analysis, reporting protocols and outbreak management" [O. Reg. 79/10, s. 229 (3)].

Infectious period –

The time during which infected individuals are able to transmit their infection to others.

Just Clean Your Hands –

A program created to help hospitals and individuals overcome the barriers to proper hand hygiene and improve compliance with hand hygiene best practices. <https://www.publichealthontario.ca/en/BrowseByTopic/InfectiousDiseases/JustCleanYourHands/Pages/Just-Clean-Your-Hands.aspx>

Kaplan's Criteria –

When microbiological confirmation of a suspected norovirus outbreak is not possible, the Kaplan Criteria may be applied to determine the likelihood that the outbreak is of viral origin. These criteria are as follows: 1) a mean (or median) illness duration of 12 to 60 hours, 2) a mean (or median) incubation period of 24 to 48 hours, 3) more than 50% of people with vomiting, and 4) no bacterial agent found. The criteria are very specific— when all four criteria are present, there is a high likelihood that the outbreak is attributable to norovirus. However, the criteria lack sensitivity—about 30% of norovirus outbreaks do not meet these criteria. Therefore, the possibility of a viral etiology should not be discarded if these criteria are not met.

Line listing –

A table that summarizes information about suspect, probable or confirmed cases associated with an outbreak. It often includes identifying information, demographics, clinical information and exposure or risk-factor information.

Long-Term Care Home (LTCH) –

A place that is licensed under the Long-Term Care Homes Act, 2007, S.O. 2007, Chapter 8, and includes a municipal home, joint home or First Nations home approved under Part VIII of the Act.

Non-staff –

Visitors, volunteers, family members and community groups.

Onset –

The date and time when clinical signs or symptoms first appear.

Outbreak –

An unexpected increase of disease occurring within a specific population at a given time and place.

Paid Sitter –

A person who is not employed by the LTCH but may provide resident care. This person is normally hired by a resident's family.

Person-to-person outbreak –

An outbreak that occurs when infection is spread from one person to another.

Point-source outbreak –

An outbreak that occurs when infections stem from a single source, for example an outbreak spread to people who have eaten a contaminated food item.

Potentially hazardous foods –

Are those that are capable of sustaining growth of pathogens (harmful bacteria). They generally have high protein content, are neutral in acidity and are moist. Common examples are poultry, meat, fish and dairy products. Potentially hazardous foods must be handled carefully with respect to temperature; they must be stored at or below 4°C and must be cooked to a specified internal temperature before being served to ensure they are safe. Although some fruits and vegetables have been implicated in large-scale, food-borne outbreaks, they lack the ingredients necessary for uncontrolled bacterial growth. It is not necessary to store them in a refrigerator or cook them to a specific temperature to ensure safety. These foods still need to be handled with extra care and washed thoroughly before use.

Resident –

An individual who permanently or temporarily lives in a LTCH.

Routine Practices –

The system of infection prevention and control practices recommended by the Public Health Agency of Canada to be used with all clients/patients/residents during all care to prevent and control the transmission of micro-organisms in health-care settings. The full description of Routine Practices to prevent and control transmission of nosocomial pathogens can be found in the Updated Best Practices Manual for Routine Practices and Additional Precautions in Ontario, available from the Public Health Ontario website at:

https://www.publichealthontario.ca/en/BrowseByTopic/InfectiousDiseases/PIDAC/Pages/Routine_Practices_Additional_Precautions.aspx

Staff –

All persons who carry on activities in the LTCH, including but not limited to employees (permanent, temporary), students, attending physicians and both health care and non- health care contract workers and any other staff (including persons with admitting/clinic privileges [MD, Mid-wives, Hearing Aid Centre]; maintenance workers (e.g. janitorial, repair, etc.) or other workers who carry on activities in resident care areas or come into contact with residents (e.g. hairdressers)].

Surveillance –

The systematic and ongoing collection, collation and analysis of data and the timely dissemination of information so that appropriate action can be taken to reduce the number of illnesses.

Visitors –

People not affiliated with the LTCH who visit residents. They may include residents' friends or relatives, or groups who visit many residents.

Volunteer –

A person who is part of the organized volunteer program of the LTCH, but are not directly employed by it (i.e they do not receive a paycheck from the LTCH).

1 Introduction

Gastroenteritis outbreaks continue to represent a significant burden of illness in Ontario's Long-Term Care Homes (LTCHs). Each year, approximately 1000 - 1200 gastroenteritis outbreaks are reported to public health. The number of cases associated with a single outbreak may range from 10 to 200. Significant mortality may also be associated with these outbreaks.

Gastroenteritis outbreaks can be caused by bacteria, viruses, or parasites contracted through the consumption of contaminated foods or beverages, and through contact with contaminated items or infected persons. Many outbreaks can be prevented or have their impact mitigated through intentional, knowledgeable and rapid identification and management of the case to minimize the spread of disease to prevent illness, hospitalization and death.

1.1 Purpose

This document will provide:

1. The minimum standards to reduce and manage gastroenteritis outbreaks in LTCHs (Box 1).
2. A list of further resources and practical tools.
3. The most updated review of evidence-based literature, clinical knowledge, trends and expert consensus on the control and management of gastroenteritis outbreaks.

Box 1. Outbreak Recommendations: Purpose and Scope

Outbreak Recommendations: Purpose and Scope

This document will provide useful and practical information that will help the LTCHs :

- Develop outbreak management policies and procedures.
- Prevent outbreaks.
- Identify and isolate cases to prevent disease spread.
- Respond to and manage a gastroenteritis outbreak should one occur.

Scope:

- May be useful in other settings – such as retirement homes, child-care, acute-care, and correctional facilities, shelters, residential camps and group homes – although not all the information in this document will be applicable to all settings.
- Replace *Control of Gastroenteritis Outbreaks in Long-Term Care Homes* (MOHLTC, 2013)
- Should be used in conjunction with other provincial and national best-practice guidelines or other guidance documents related to gastroenteritis and infection prevention and control. See References and Further Information Section.

1.2 The Role of Public Health

Local boards of health provide invaluable assistance and expertise in the prevention, detection, management, and control of gastroenteritis outbreaks. Health-care facilities and institutions are required to report gastroenteritis outbreaks to the Medical Officer of Health. The Ontario Public Health Standards: Requirements for Programs, Services, and Accountability state that each board of health shall work with community partners and service providers to determine and address the need for knowledge translation resources and supports in the area of infection prevention and control; and the board of health shall work with the appropriate partners to increase awareness among relevant community partners, including correctional facilities, health care, and other service providers, of: the local epidemiology of communicable disease and other infectious disease of public health importance; infection prevention and control practices; and reporting requirements for diseases of public health significance, as specified in the *Health Protection and Promotion Act*.

The roles and responsibilities of the Public Health Unit with respect to gastroenteritis outbreaks are outlined in Box 2.

Box 2. Public Health Units: Roles and Responsibilities

Public Health Units: Roles and Responsibilities
<ul style="list-style-type: none">• Ensuring qualified public health personnel are available 24 hours a day, 7 days a week.• Supporting/consulting with Infection Prevention and Control Professionals and providing representation on the OMT.• Supporting Infection prevention and control programs and committees.• Consulting on surveillance, infection prevention and control policies and procedures, and outbreak management.• Consulting on the investigation and management of communicable and reportable diseases.• Collecting data, analyzing outbreaks, and reporting to the MOHLTC on institutional gastroenteritis outbreaks for use in further analysis and the identification of provincial trends.• Performing regular food and environmental inspections.• Food and equipment testing may be required during the investigation of food-borne outbreaks.• Ensuring LTCH staff has ready access to the contact names/numbers for the Public Health Unit. <p>http://www.health.gov.on.ca/en/common/system/services/phu/locations.aspx</p>

1.3 The Role and Responsibilities of the Long-Term Care Homes

LTCHs are responsible for ensuring effective outbreak management, timely detection of cases and communication of these cases to public health. The requirement for communication to public health is stipulated under Section 27(2) of the HPPA. LTCHs are also responsible for reporting outbreaks of reportable or communicable disease as defined in the HPPA to the Director named in the LTCHA (O. Reg. 79/10, s.107(1)5.). Infection control measures recommended for all health care settings and LTCHs are outlined in *Best practices for infection prevention and control programs in Ontario* (Appendix B in PIDAC 2012).

1.4 Types of Gastroenteritis Outbreaks

Gastroenteritis outbreaks may result from person-to-person spread or ingestion of contaminated food or water. Contamination of food or water may occur at the source or during their preparation, handling or storage. Indirect transmission may occur through contact with contaminated fomites. The agents responsible for these outbreaks may be viral, bacterial or parasitic in nature.

1.4.1 Outbreaks Caused by Viruses

Viral gastroenteritis is the leading cause of gastroenteritis outbreaks in institutions. In LTCHs, norovirus is the most common cause of such outbreaks. The modes of transmission for norovirus include aerosolization, indirect transmission via contaminated surfaces, person-to-person spread or consumption of contaminated food and beverages. It is of importance to note that contamination of food most often occurs by an infected food handler. Norovirus affects both residents and staff, especially during the winter months when community incidence is also high. Indicators of a norovirus outbreak include sudden onset of symptoms with a significant proportion of affected persons experiencing nausea, vomiting and diarrhea. Norovirus outbreaks in LTCHs can place increased stress on resources, including increased costs due to higher demands for health care workers (Zingg et al. 2005). During a viral outbreak, more than 50% of the residents and staff may become ill. Factors involved in viral outbreaks are outlined in Box 3.

Although there may be a common source such as a food item that is responsible for an outbreak, secondary transmission from person-to-person can readily occur. Infected

individuals typically shed millions of viral particles in their feces or vomitus. However, only a few of these particles are needed to cause infection. Norovirus particles can contaminate large portions of the environment in a LTCH and are able to survive for days on a variety of surfaces, making hand hygiene vital to infection prevention and control efforts.

Box 3. Examples of How a Viral Outbreak Starts

Examples of How a Viral Outbreak Starts?
<ul style="list-style-type: none">• An infected person with unclear hands or contaminating commonly-touched surfaces and equipment• An infected staff or family member with unclean hands or gloves providing care to residents.• Inadequate cleanup of body fluids (i.e., diarrhea and vomit) and the subsequent contamination of the environment.• Vomiting and/or uncontrolled diarrhea spreading virus, through droplets sprayed into the air (aerosolization).• The sharing of resident equipment, such as a commode, that is not properly cleaned and disinfected between uses by different residents.• Any person who enters the LTCH when they have symptoms of gastroenteritis.• An infected food handler contaminating residents' food.

1.4.2 Outbreaks Caused by Bacteria and Parasites

Bacteria and parasites are less frequently implicated in gastroenteritis outbreaks. Such outbreaks often arise from a point source such as bacteria-contaminated food or water. The initial attack rate can be high, but the disease usually does not spread beyond those initially infected. Unlike viral transmission, person-to-person transmission of bacteria and parasites is less common. As a result, there is greater success in controlling outbreaks caused by bacteria and parasites. Some bacterial pathogens such as *Shigella*, Verotoxin-producing *Escherichia coli* and *Salmonella typhi* require special considerations. LTCHs should consult with the public health unit for further guidance.

For specific infection prevention and control practice information on *Clostridium difficile* Infection (CDI), please see the PIDAC document, Annex C: *Testing, Surveillance and Management of Clostridium difficile in All Health Care Settings* (PIDAC – 2013).

For information on the roles and responsibilities associated with the reporting and

management of CDI outbreaks, please see the MOHLTC document Roles and Responsibilities of Hospitals and Public Health Units for Clostridium difficile Infection (CDI) Reporting and Outbreak Management (MOHLTC – 2013)

2 Preparation – Infection Control and Outbreak Policies and Procedures

Not all gastroenteritis infections and outbreaks in LTCHs are preventable. However, 20% of these infections can be prevented through adherence to an Infection Prevention and Control (IPAC) program (Harbath et al. 2003).

While infection prevention and control is everyone's responsibility, a staff member with education and experience in IPAC must be designated to co-ordinate the program. An alternate should also be assigned during the IPAC program coordinator's absence.

Effective IPAC begins with preparation and implementation of outbreak control policies and procedures, the use of infection prevention techniques (Routine Practices) and the implementation of Additional Precautions when deemed necessary.

Recommendations for developing outbreak control policies and procedures are outlined in Box 4.

The following documents are also recommended for this purpose:

- *Best practices for infection prevention and control programs in Ontario* (PIDAC 2012).
- *Best practices for Environmental Cleaning for Prevention and Control of Infections* (PIDAC 2012).
- *Infection control guidelines: Routine Practices and Additional Precautions for Preventing the Transmission of Infection in Health Care* (PHAC 1999).
- *Routine Practices and Additional Precautions in all Health Care Settings* (PIDAC 2012).

Box 4. Policy and Procedure Preparation

Policy and Procedure Preparation

- **Disease Prevention:** include Routine Practices and Additional Precautions; hand hygiene; reducing the risk of transmission of infectious agents; education of staff, volunteers, residents and families; and environmental cleaning.
- **Outbreak Preparedness:** policies should be evaluated and updated at least annually in accordance with evidence-based practices or prevailing practices.
- **Surveillance:** include early identification of cases and application of appropriate practices to prevent disease transmission, management of data and the declaration of an outbreak.
- **Reporting:** Include notification of the public health unit and other relevant authorities
- **Management of an Outbreak:** include composition, mandate and roles of the Outbreak Management Team (OMT); a policy on staff exclusion; an outbreak staffing/resident plan for cohorting, transferring and workload management to ensure an adequate staff-to-resident ratio.
- **Staff and Volunteer Policies:** provide orientation to hand hygiene; modes of transmission; cleaning and disinfection; and use of personal protective equipment. Retraining of staff should be in accordance with evidence-based practices or prevailing practices.
- **Resident and Visitor Policies:** educate on what will happen during an outbreak.

2.1 Outbreak Preparedness

As part of the IPAC program, all policies and procedures should be evaluated and updated annually by the LTCH. All LTCHs are encouraged to include Public Health in the evaluation. Every LTCH should develop policies and procedures to include:

- The composition and mandate of the OMT.
- A policy on excluding staff – and the criteria for their returning to work – during a gastroenteritis outbreak.
- An outbreak staffing/resident plan for cohorting, transferring and workload management.
- Specimen collection, including where to obtain specimen kits; testing facilities available and after hours testing contact information (Appendix 1). Policies should also address receiving and reporting of laboratory test results.

- The carrying out of control measures for residents, staff, visitors and volunteers – including education about and reinforcement of Routine Practices and Additional Precautions and other control measures as applicable (Appendices 2-7).
- Routine, thorough cleaning and education on specific disinfection procedures, depending on the identified type of organism (PIDAC 2010).
- Food safety, records such as temperature logs for food and dishwashers are recommended. Menu choices; catered and external food for residents, and routine retention of food samples should also be recorded.
- The roles and responsibilities of the home during an outbreak which include contact with the public health unit, the laboratory or laboratory-testing services available, communication with residents and their families, staff, volunteers, external groups and media during an outbreak.
- Procedures for declaring the outbreak over.

Infection Prevention and Control and Workers' Safety

When preparing for gastroenteritis outbreaks, it is often viewed as a resident safety issue; however, the health of workers in LTCHs should be considered. LTCHs must comply with applicable provisions of the *Occupational Health and Safety Act* (OHSA) and its Regulations when implementing infection prevention and control procedures since infection of staff is an occupational health and safety issue (Box 5). Employers, supervisors, and workers have rights, duties, and obligations under the Act. A proactive approach to preparing policies and procedures is required to reduce the impact of outbreaks in LTCHs (PIDAC 2011).

Staff safety is addressed in OHSA and its Regulations, including the use of personal protective equipment (PPE), regulations pertaining to the proximity of food and drink to infectious materials, needle safety, and ceiling exposure values for biological and chemical agents.

Box 5. Health and Safety of Health Care Workers

Health and Safety of Health Care Workers

The *Regulation for Health Care and Residential Facilities* under the *Occupational Health and Safety Act* (OHSA) requires LTCHs develop written measures and procedures for the health and safety of workers, in consultation with the LTCHs Joint Health and Safety Committee

These measures and procedures may deal with:

- Proper hygiene practices and the use of hygiene facilities.
- Control of infections.
- Use of appropriate antiseptics, disinfectants and decontaminants.
- Use, application, care, removal and limitations of PPE.
- Development of health and safety training and educational programs for workers by employer, that are relevant to the workers' jobs.

2.2 The Infection Prevention and Control Program – Disease Prevention

As noted previously, effective IPC programs can reduce infections in health care settings. Central to an effective IPC program is the implementation of Routine Practices and Additional Precautions. These measures should be at the core of a disease prevention culture in all LTCHs.

2.2.1 Routine Practices

Routine Practices are based on the premise that all residents are potentially infectious even when asymptomatic, and that standards of practice should be applied **routinely** with **all** residents during all care to prevent exposure to blood, body fluids, secretions, excretions, mucous membranes, non-intact skin or soiled items and to prevent the spread of micro-organisms (Box 6; Appendices 2-7). Healthcare providers must assess the risk of exposure to blood, body fluids and non-intact skin and identify the strategies that will decrease exposure risk and prevent transmission of micro-organisms before interacting with the resident.

Routine Practices for specific gastrointestinal micro-organisms and conditions are outlined in Appendix 2. Each health-care provider must conduct a risk assessment before interacting with the resident. To assess the risk of exposure, the health-care worker should consider:

- Risk of exposure to body fluids or blood during the procedure. Thorough hand hygiene is sufficient for minimal risk procedures, whereas higher risk procedures require both thorough hand hygiene and use of additional infection control practices.
- The procedure and the skill level of the health-care worker performing the procedure. Usually, the better trained a health-care worker is, the less likely they will be exposed to body fluids or blood.
- The resident's level of cooperation and cognitive awareness; for example, the more cooperative/cognitively aware, the lower the risk of transmission.
- Which infection prevention strategies to use during the resident and health-care-provider interaction.

Box 6. What are Routine Practices?

What are Routine Practices

Routine Practices is the system of infection prevention and control practices recommended by the Public Health Agency of Canada to be used with all residents during all care, to prevent and control the transmission of micro-organisms in health-care settings. *Routine Practices fact sheet* is available at:

https://www.publichealthontario.ca/en/BrowseByTopic/InfectiousDiseases/PIDAC/Pages/Routine_Practices_Additional_Precautions.aspx

These practices describe prevention and control strategies to be used with all residents during all resident care, and include:

- Hand hygiene with an alcohol-based hand rub or with soap and water before and after any direct contact with a resident.
- The use of additional barrier precautions to prevent staff contact with a resident's blood, body fluids, secretions, excretions, non-intact skin or mucous membranes.
- Gloves are to be worn when there is a risk of hand contact with a resident's blood, body fluids, secretions, excretions, non-intact skin or mucous membranes; gloves should be used as an additional measure, not as a substitute for hand hygiene.
- Gowns are to be worn if contamination of uniform or clothing is anticipated.
- The wearing of masks and eye protection or face shields where appropriate, to protect the mucous membranes of the eyes, nose and mouth during procedures and resident care activities likely to generate splashes or sprays of blood, body fluids, secretions or excretions.

Hand Hygiene

Proper hand hygiene is the single most important practice in preventing the transmission of infections. Hand hygiene refers to any action of cleaning one's hands; this may involve using alcohol-based hand rub or liquid soap and water. Hand hygiene also encompasses the use of skin conditioners to maintain skin integrity, keeping nails short and clean, and refraining from wearing jewelry or nail enhancements.

To remove and/or kill micro-organisms on hands:

- Wash with soap and running water.
- When hands are not visibly soiled, use an alcohol-based hand rub (ABHR) containing at least 70% alcohol.

LTCHs should emphasize the importance of proper hand hygiene through ongoing education and regular communications with staff, residents, visitors and volunteers (Appendix 5). Auditing hand hygiene practices has been shown to improve compliance. Audit performance of hand-hygiene to identify current practices and to subsequently develop and implement strategies for improving thoroughness, appropriate timing and increased frequency. These audits should include direct observation.

Box 7. Hand Hygiene Program

Hand Hygiene Program

Essential Components of the Hand Hygiene Program:

- Develop a policy and procedure for hand hygiene and hand care programs.
- Ensure easy access to hand hygiene agents and to hand washing sinks dedicated to hand hygiene and used for no other purpose.
- 70 to 90% alcohol-based hand rubs must be provided in the LTCH.
- Education that includes indications for hand hygiene, techniques, indications for hand hygiene agents and hand care.
- Program to monitor hand hygiene compliance with feedback to individuals and management via the Infection Prevention and Control Committee.
- Just Clean Your Hands material is available for Long-Term Care Homes at <https://www.publichealthontario.ca/en/BrowseByTopic/InfectiousDiseases/JustCleanYourHands/Pages/Just-Clean-Your-Hands.aspx>

Personal Protective Equipment (PPE)

PPE, such as gloves and gowns, and in some situations masks and eye or face protection, may be required as barriers against micro-organisms. When used correctly, PPE helps protect staff and residents from infections and the environment from contamination. PPE only works when used properly. It is important for staff to don and doff PPE properly to avoid contaminating themselves, residents and the environment (Appendices 6, 7; Section 2.3.1 Contact Precautions).

Gloves

Gloves protect the hands of health-care providers from contact with the resident's body fluids, blood, excretions, secretions or non-intact skin. The **do's** and **don'ts** of using gloves are outlined in Box 8. **Note:** Not all gloves are suitable for all tasks. For example, co-polymer gloves are not suitable for providing direct care to residents, but may be used in food preparation. Match the type of glove to the procedure and the strain exerted on the glove during the procedure. The ICP, public health unit or Regional Infection Control Network (RICN) can help in choosing appropriate gloves. Be aware of possible latex allergies, latex gloves are rarely necessary and should not be used if there is risk that staff or residents may have a latex allergy. A comparative document to medical-grade gloves is presented in *Routine practices and additional precautions in all health care settings* (Appendix 6).

Gowns

Use gowns to protect uncovered skin and protect clothing or uniforms during activities likely to generate splashes or sprays of blood, body fluids, secretions or excretions. The cuffs of the gloves must cover the cuffs of the gown. Remove gowns before leaving the resident's room or dedicated space or if they become soiled, wet or contaminated.

Masks and Eye Protection

Wear masks and eye protection to protect eyes, nose and mouth during procedures likely to generate splashes or sprays of blood, body fluids, secretions or excretions. Consider using a mask to minimize the risk of touching nose and mouth with contaminated hands.

Box 8. Effective Use of Gloves for Disease Prevention

Effective Use of Gloves for Disease Prevention

DO NOT:

- Substitute gloves for hand hygiene.
- Wear gloves for routine procedures limited to touching a resident's intact skin.
- Wear gloves when feeding a resident.
- Wash or reuse single-use, disposable gloves.
- Double glove.
- Wear the same gloves for activities involving more than one resident.

DO:

- Perform hand hygiene and wear clean, non-sterile gloves when:
 - i. Contact blood, body secretions or excretions is anticipated.
 - ii. Handling visibly soiled items.
 - iii. Staff's hands have open cuts, wounds or skin conditions i.e. eczema, psoriasis, dermatitis.
- Change gloves and perform hand hygiene before putting on new gloves and between procedures with the same resident.
- Remove and discard gloves:
 - i. Immediately after completion of the task.
 - ii. If they become ripped or damaged.
 - iii. At the point of use before leaving the room.

2.3 Additional Precautions

Additional Precautions are necessary for certain pathogens and symptoms, and includes airborne, droplet and contact modes of transmission (PIDAC 2012).

Gastrointestinal infections typically cause significant vomiting and diarrhea, which can contaminate the environment. Contact and/or droplet precautions are recommended as soon as symptoms of a gastroenteritis infection develop as the virus may become aerosolized.

2.3.1 Contact Precautions

Contact Precautions should always be used in addition to Routine Practices. The following strategies will help decrease transmission during a gastroenteritis outbreak (see Appendices 2, 4):

- Encourage residents with gastrointestinal symptoms to remain in their rooms and provide them with tray service.
- Do not allow infected residents to participate in group activities for at least 48 hours after their symptoms have resolved.
- Instruct visitors on the precautions they should follow.
- Identify, and store separately, all equipment designated to be used by an ill resident, to prevent their use for other residents. If a lack of equipment or storage space makes this unfeasible, then do not use the equipment until it has been thoroughly cleaned and disinfected.
- Wear gloves and gown when providing direct care to a resident.
- Wear a mask and goggles or a face shield to protect from splashes if a resident has explosive diarrhea or projectile vomiting or when there are other situations that pose risk of splashing such as removing fecal material or vomitus, or using a sink spray nozzle. To prevent contamination, store clean supplies outside the rooms of infected residents.
- Provide containers in residents' rooms for used PPE disposal.
- Reinforce the importance of hand hygiene with roommates and visitors – visitors who provide direct care to residents (i.e. toileting) should use the same PPE as staff and be instructed on how to properly do so.

2.3.2 Other Disease Prevention Strategies

Other disease prevention strategies include, routine environmental cleaning and disinfection, safe food-handling practices, a staff health policy and a visitor health policy, which shall include but not be limited to the recommendations below (PIDAC 2012).

Environmental Cleaning and Disinfection:

Each LTCH should have written policies and procedures for:

- Routine cleaning and disinfecting.
- Enhanced cleaning during an outbreak. Enhanced environmental cleaning practices should always be implemented during outbreaks (Box 9).

These policies and procedures should be evaluated and updated at least annually to ensure they reflect current best practices. Policies should include proper use of supplies for cleaning and disinfecting; laundry-handling practices; and proper handling and disposal of waste (Appendix 8, PHAC 1998, PIDAC 2011, PIDAC 2012).

Some basic cleaning principles include:

- Moving from clean areas to dirty areas.
- Cleaning from top to bottom.
- Increased cleaning of high-touch surfaces during gastroenteritis outbreaks.
- Adhering to manufacturer's instructions on preparation and storage of disinfectant solutions and the recommended contact time.

Box 9. Enhanced Environmental Cleaning Practices

Enhanced Environmental Cleaning Practices
<p>Emphasize the following during an outbreak:</p> <ul style="list-style-type: none">• Increase routine cleaning of all high-touch surfaces such as door handles, bed railings, hand rails, light switches, elevator buttons, over-bed tables, dining tables and counters.• Increase the cleaning and disinfecting of all surfaces in the ill resident's immediate environment.• Disinfect shared resident equipment after each use and discard disposable equipment before leaving the resident's room.• If possible, dedicate specific equipment to each ill resident.• Promptly clean and disinfect surfaces contaminated by stool and vomit (Appendix 8).• Clean soiled carpets and soft furnishings with hot water and detergent or steam clean – vacuum cleaning is not recommended.

Safe Food Handling Practices:

The Public Health Inspector can provide information on food safety requirements for:

- Proper food-handling techniques.
- Food-service worker hygiene.
- Sanitation.
- Food-safety education.

Staff Health Policy: Staff with any gastroenteritis symptoms should stay off work when ill and continue to remain home for a minimum of 48 hours after their GI symptoms (i.e. vomiting, diarrhea) have resolved. The LTCH can consult with Public Health once a specific agent has been identified where the exclusion period may need to be adjusted.

Visitor Health Policy: Policies must be in place for visitors that are either infectious or who may become infected. Some policies that will help protect the residents, staff and visitors include:

- Posting signs at entrances outlining screening policies for non-staff (Appendix 9).
- Advising everyone not to enter the LTCH when they have gastrointestinal symptoms, respiratory symptoms or known communicable disease. Visitors should not enter the LTCH if they have symptoms of a communicable disease, particularly if they are experiencing diarrhea or vomiting. This applies at all times, not only during outbreaks.
- Providing hand-hygiene products at the entrances and throughout the LTCH.

2.3.3 Education of Staff and Volunteers

An effective education program that is well planned and executed will improve IPAC programs (Daly et al. 1992). The LTCH should educate all staff and volunteers about gastrointestinal infections at the time of hiring or orientation, annually thereafter, and when an outbreak occurs (Box 10). A mechanism to track all education should be implemented. Education can include the use of brochures, signs and posters as well as courses and demonstrations. The type of education provided should be tailored to meet the needs of staff and volunteers and the specific activities that they carry out within the LTCH. For further information or assistance, contact your public health unit or RICN office.

Box 10. Education for all Staff and Volunteers

Education for all Staff and Volunteers

Education/orientation programs for all staff and volunteers (as applicable) should include information on:

- The transmission and prevention of gastrointestinal infections.
- Routine Practices (Box 5, Appendices 2-7) and Additional Precautions (Section 2.3).
- Hand hygiene (Appendix 5).
- Donning and doffing PPE (Appendices 6, 7).
- Appropriate cleaning and disinfecting procedures to be followed after each use of multi- use equipment which is shared among residents (Appendix 8).
- Environmental cleaning and disinfecting procedures, especially for housekeeping staff.
- Food safety – safe food-handling practices for receiving, preparing, storing and transporting food.
- Occupational Health policies.
- Routine daily surveillance for signs of infection.
- Roles and responsibilities of staff, administration, the ICP and Public Health.
- Specimen collection methods (Appendix 1).
- Gastroenteritis outbreak management.
- Outbreak control.

2.3.4 Education of Residents and Visitors

Provide residents and visitors with appropriate education on:

- Hand hygiene.
- Routine Practices and, in specific instances, Additional Precautions (specifically proper use of PPE).
- The transmission and prevention of gastrointestinal disease including the requirement not to visit the LTCH when they are ill.
- Outbreak Management – what to expect (such as restrictions while visiting) when there is a gastroenteritis outbreak.

3 Surveillance and Outbreak Detection

3.1 Surveillance

Effective surveillance is essential to ensure the early identification of outbreaks so that control measures can be instituted as soon as possible. LTCHs are required to have an ongoing surveillance program to detect the presence of infections in the LTCH.

Surveillance is the systematic ongoing collection, collation and analysis of data with timely communication and sharing of information in order to ensure that disease aberrations and outbreaks are responded in an appropriate and effective manner (Box 11).

3.1.1 Target Groups for Surveillance

Resident Surveillance

Direct-care staff members are key to good resident surveillance; however, all staff should be encouraged to report changes in any resident condition. Staff who recognize and report initial signs of resident illness allow control measures to be implemented early, which is a vital step in preventing an outbreak.

To provide effective surveillance, staff must be given education and/or training in:

- Their role in surveillance and its importance.
- Symptoms of gastrointestinal infection.
- Criteria for a suspected outbreak.
- Procedures for reporting to the ICP or designate, who will then report to the public health unit.

LTCHs must identify residents with gastrointestinal symptoms on a daily surveillance form or line list (Box 11, Appendix 10). Completed surveillance forms should be forwarded to the LTCHs ICP or designate daily. Typical surveillance strategies include:

- Conducting unit rounds.
- Reviewing unit reports that include incidents of diarrhea and vomiting.
- Reviewing physician/staff communication books.
- Reviewing medical and/or nursing progress notes in charts.
- Reviewing pharmacy utilization records.

- Reviewing laboratory reports.
- Receiving verbal reports from unit staff based on their clinical observations.

Box 11. The Surveillance Form

The Surveillance Form
<p>What to include in your Surveillance Form:</p> <ul style="list-style-type: none"> • Resident name and location in LTCH. • Gender and age or date of birth. • Signs and symptoms related to the gastrointestinal infection. • Onset date of symptom(s). • Diagnostic tests and results when available. • Recent history of movement inside and outside the LTCH, such as hospitalization, diagnostic testing, clinical assessments and outings. • Underlying conditions or medications, such as laxative and antibiotic use that may cause gastrointestinal symptoms. • Sample form, Appendix 10.

Staff Surveillance

The ICP should monitor staff illness to identify clusters or outbreaks. Since ill staff can bring infectious diseases into the LTCH, staff should:

- Self-screen and stay home when they are vomiting and/or have diarrhea.
- Report vomiting and/or diarrhea to their supervisor or a person accountable for employee health. The Occupational Health designate must promptly inform the ICP of cases and clusters of staff who are absent from work with the symptoms of gastroenteritis.

Analysis

The ICP or designate must review surveillance data daily to determine whether any resident has symptoms of infectious gastroenteritis and if more than one person has such symptoms (i.e. an outbreak is suspected, Section 3.2). The public health unit can help interpret and analyze surveillance data (See Appendix 10).

3.2 Outbreak Detection

The case definitions for a gastroenteritis outbreak use standard symptoms. These definitions apply in most situations but may need modification as circumstances dictate.

The case definition contains the criteria to be used during an outbreak to designate a

resident or staff member as having infectious gastroenteritis. The case definition developed for residents may be different from that for staff. Individuals who meet the case definition are considered a case – even if laboratory test results are negative – unless another diagnosis or reason for symptoms (e.g., laxative use) is confirmed or the case definition is changed to include the laboratory diagnosis.

Note: The Medical Officer of Health or designate may declare an outbreak in situations where **new, unusual or emergent diseases** may be suspected or identified.

3.2.1 Infectious Gastroenteritis Case Definition

To be defined as a case of infectious gastroenteritis, **at least one** of the following must be met:

- Two or more episodes of diarrhea or watery stool (takes the form of its container) within a 24-hour period, or two or more episodes of vomiting within a 24-hour period, or
- One episode of diarrhea or watery stool (takes the form of its container) and one episode of vomiting within a 24-hour period, or
- Laboratory confirmation of a known gastrointestinal pathogen and at least one symptom compatible with gastrointestinal infection (e.g., nausea, vomiting, diarrhea, or abdominal pain or tenderness)

Note: Care should be taken to rule out non-infectious causes of these symptoms such as new medications, use of laxatives or other non-infectious diseases. The bowel movements should be unusual or different for the resident. Some residents may not be able to report nausea or abdominal pain, in which case careful observation may be needed to determine if these symptoms are occurring. For example, behavioural changes may be a clue that residents are experiencing nausea or pain. Frail residents with small appetites may have only one episode of either vomiting or diarrhea and may or may not exhibit other signs and symptoms associated with gastrointestinal illness. LTCHs may want to consider these residents as suspect cases and implement infection-control measures to prevent potential transmission.

3.2.2 Outbreak Definition

The LTCH should establish a baseline of normal occurrences of gastroenteritis. A gastroenteritis outbreak is defined as the occurrence of gastroenteritis beyond what is normally expected based on surveillance data. This definition makes it clear why it is important to conduct ongoing surveillance. Additional outbreak definitions are provided below for guidance. Contact your Public Health Unit immediately if you suspect an outbreak.

3.2.3 Suspected Gastroenteritis Outbreak Definition

Two suspected cases of infectious gastroenteritis in a specific area, such as a home, unit, or floor within 48 hours.

3.2.4 Gastroenteritis Outbreak Definition

Three or more cases of infectious gastroenteritis in a specific area within a four-day period, or three or more units/floors having a case of infectious gastroenteritis within 48 hours. **Note:** This definition may be modified as the investigation proceeds.

4 Managing Gastroenteritis Outbreaks

Even a relatively small gastroenteritis outbreak is disruptive, often frustrating and exhausting for everyone involved. Infection control efforts can be difficult even with consistent use of Routine Practices (Appendices 2-7) and other outbreak control procedures, however, early detection and management are the keys to minimizing the impact. As a quick reference document, the steps involved in managing a gastroenteritis outbreak are presented in Appendices 11 and 12. Your public health unit will also have outbreak management resources to help you prepare.

4.1 Control Measures for a Suspected Gastroenteritis Outbreak

In the event of a suspected outbreak, the LTCH should immediately institute control measures, including, but not limited to:

- Advising Public Health.
- Encouraging residents with suspected gastroenteritis to remain in their rooms and providing them with tray food service.
- Reviewing Routine Practices.
- Assessing the need for Contact Precautions (Section 2.3.1).
- Enhancing surveillance to look for additional residents and staff that meet the case definition.
- Reinforcing the importance of proper hand hygiene and other Routine Practices.

4.2 Steps in Outbreak Management

Each LTCH is responsible for ensuring that these outbreak management steps are implemented. Many of these activities are done simultaneously. **Use Appendix 11 and 12 as a quick reference** for steps to take during an outbreak.

4.2.1 Assess the Outbreak

A line-listing template is available from your Public Health Unit. The template provides documentation of the extent and nature of the suspected outbreak (Appendix 13). Begin a line listing by adding surveillance data from the daily sheets. Prepare separate line lists for residents and staff and, if useful, keep a separate line listing for each affected unit/floor. The line listing may be expanded to include other relevant data – beyond that recommended in this document – as the investigation proceeds. Identify those at highest risk of infection within the LTCH.

4.2.2 Implement General Infection Prevention and Control Measures

Implement control measures and notify all staff as soon as an outbreak is suspected (See Section 5.0). Implement enhanced environmental cleaning (Refer to Box 9, section 2.3.2.). For further information or assistance, contact your public health unit or RICN office.

4.2.3 Consult with the Public Health Unit

Upon notifying the Public Health Unit, provide an updated line listing, the name of the primary contact person responsible for the outbreak investigation (usually the ICP) and the names of staff responsible for managing the outbreak on evenings, weekends, holidays and during vacation periods. Follow the remaining steps:

- Review the preliminary case definition for the suspected outbreak. Confirm with the Public Health Unit that infection prevention and control measures have been implemented.
- Request an Outbreak Number for the investigation.
- Discuss specimen collection and testing procedures with your Public Health Unit to determine:
 - The number of specimens that will be collected, stored and submitted to the laboratory.
 - The number of laboratory specimens that should be taken during the initial outbreak investigation.
 - Which residents to test during the initial outbreak investigation.
 - How the specimens should be handled (e.g. stored, transported to the lab).

4.2.4 Declare an Outbreak

Outbreaks are declared by the Medical Officer of Health (or their designate), the Medical Director of the LTCH or the Director of Nursing and Personal Care of the LTCH based on either:

- The definition in this document; or
- Other specific factors of the particular outbreak.

Case definitions should be reviewed periodically during the course of the outbreak and modified if necessary, for example, when you identify residents with new symptoms.

4.2.5 Notify Appropriate Individuals/Agencies

In addition to the Public Health Unit, notify some or all of the following as appropriate, both internally and externally.

INTERNALLY:

- Medical Director
- Director of Nursing and Personal Care
- Administrator
- Licensee and/or Board of Directors
- Chair of the Infection Prevention and Control Committee, Infection Prevention and Control Professional
- Employee Health Nurse
- Families of all residents
- Director of Food Services
- Director of Housekeeping/Maintenance
- Resident and Family Councils
- All Residents
- All Staff members
- All Volunteers
- Attending physician(s), Nurse Practitioners, Resident physicians
- Other health-care providers, such as physiotherapists
- Other service providers, such as hairdresser

EXTERNALLY:

- Ministry of Health and Long-Term Care through the Critical Incident System
- Acute-care hospitals – Infection Prevention and Control Professional
- Admitting and Emergency departments

- Community Care Access Center /other institutions
- Staffing agencies
- Emergency Medical Services – ambulance
- Coroner’s office if there is a death immediately preceding/during the outbreak
- Ontario Ministry of Labour, Regional Office (for cases of health-care acquired gastroenteritis in staff members)
- Laboratory Services Provider
- Pharmacist/Advising Pharmacy
- Local Health Integration Network (LHIN)
- The Provincial Transfer Authorization Centre (PTAC)
- Support services

4.2.6 Call an Outbreak Management Team Meeting

Administration of the LTCH should hold a meeting with LTCH representatives from each department who have decision-making authority and a Public Health Unit representative as soon as possible. This group will become the Outbreak Management Team (OMT) that meets daily to manage all aspects of the outbreak (Box 12).

Outbreak Management Team Duties:

Surveillance

- Review line-listing information to ensure OMT members understand the current status of the outbreak, its progression and the population at risk.
- Review case and outbreak definitions to ensure all members of the OMT have a common understanding of the surveillance criteria.

Investigation

- Confirm arrangements for the collection/submission of specimens for laboratory analysis (Appendix 1).
- Work with your Public Health Unit, RICN and other experts as needed.

Implementation of Control Measures

- Review the control measures and recommend any necessary modifications.
- Control measures may include:
 - Enhanced hand hygiene.
 - Strict adherence to Routine Practices and Additional Precautions.
 - Enhanced environmental cleaning.
 - Enhanced visitor monitoring.
 - Restricting activities.
 - Modified food service operations if food is relevant to the outbreak.
- Confirm that the LTCHs ICP, or designate, is responsible for ensuring the agreed upon control measures are in place and enforced.

Communications

- Review who (persons/institutions) have been notified of the outbreak.
- Prepare/distribute internal communications for residents, families, staff and volunteers.
- Review process for ongoing communication/education for staff including the individual who is responsible.
- Review the process for communicating laboratory results and control measures with Public Health Unit staff and the ICP.
- Confirm the process for daily communication between the LTCH and the Public Health Unit.
- Ensure that the Public Health Unit and the LTCHs contact telephone numbers are accurate/available.
- Verify contact information for after-hours, weekends and holidays.
- Confirm appropriate outbreak notification signage is available and posted.

Box 12. The Outbreak Management Team (OMT)

The Outbreak Management Team (OMT)
<p>Roles and Responsibilities:</p> <p>Chairperson</p> <ul style="list-style-type: none">• Coordinates outbreak control meetings and the agenda.• Delegates tasks. <p>Outbreak Coordinator (often the ICP)</p> <ul style="list-style-type: none">• Ensures all OMT decisions are carried out.• Coordinates all activities required to investigate/manage the outbreak. <p>Secretary</p> <ul style="list-style-type: none">• Sets meeting times, location, and notifies committee members of any changes.• Records and distributes minutes of meetings. <p>Media Spokesperson</p> <ul style="list-style-type: none">• Only the representative(s) identified by the OMT as the spokesperson(s) should give information to the news media. The media spokesperson can be a representative from the LTCH, the Public Health Unit or a representative from each organization.

Communications' Role with the Media

- Confirm that the Public Health Unit may release information (including the name of the LTCH) to the media or others as necessary for transparency and to reduce the risk of disease in the community or other facilities within the Public Health Unit's jurisdiction.
- Develop a crisis communications plan in advance.
- Prepare an outbreak-specific communication plan.
- Prepare a media release if appropriate and confirm media spokesperson(s).

Evaluation

- Develop an evaluation framework for use during the outbreak and after it has been declared over. The evaluation framework should include the effectiveness of outbreak control measures throughout the course of the outbreak.

4.2.7 Monitor the Outbreak on an Ongoing Basis

Outbreak monitoring should include:

- Ongoing surveillance to identify new cases.
- Monitoring the status of ill residents and staff.
- Updating line listings (Box 13).
- Ongoing monitoring of precautions and control measures.
- Reporting any significant changes in the nature of the outbreak (e.g. hospitalizations, deaths, changes in clinical picture).

The ICP must update the line listing and inform the Public Health Unit contact on a daily basis or as previously arranged. The review of the updated information should examine if there has been ongoing transmission, the effectiveness of control measures and the presentation of the new cases. Furthermore, the review may suggest changes in the outbreak definition, changes in outbreak control measures and additional laboratory testing (keeping in mind the possibility that additional causative micro-organisms could be present).

5 Control Measures For Specific Groups

5.1 Control Measures for Residents

The recommendations contained in this document are intended to protect the health of the resident populations. Recommendations are made in the interest of the resident populations at risk and as a result have the potential to affect a LTCH licensee's obligation to fully respect and promote individual resident rights as set out in the Bill of Rights under the *Long-Term Care Homes Act, 2007*, which came into force on July 1, 2010. The LTCH and PHU should work together to reduce the risk of non-compliance with the LTCHA and the HPPA, ensure that residents' rights under the LTCHA are fully respected and promoted and ensure that outbreak control measures are appropriate and proportional to the risk profile of the outbreak.

Consideration for *individual resident's* rights has always been important to PHU staff when providing outbreak management recommendations; however, public health units, under the authority of the HPPA, make recommendations aimed at protecting the health of *resident populations* in LTCHs.

When communicating outbreak control measures and recommendations to the LTCH, public health unit staff will need to emphasize the need for adherence to IPC principles with respect to exceptional visit requests; LTCH staff should be advised to call the public health unit on how to proceed, if there are any concerns regarding how to mitigate the infection control risks of a particular request from a resident/resident's family. Examples include a request for allowing children to visit during an outbreak because they don't have child sized PPE or if a visitor wishes to visit numerous residents.

The LTCH infection control practitioner or the most responsible person should contact the public health unit in order to balance the needs of the resident against the risk to the health of the other residents; at this point, a discussion around if/how the request can be accommodated can take place.

When providing outbreak management recommendations, public health units will have to assess the risk of non-compliance to outbreak control measures on the general resident population.

Generally, LTCHs and PHUs discuss with OMT members the gastroenteric infection outbreak control measures and decide jointly on appropriate measures to implement. The extent to which outbreak control measures can be implemented and what is considered reasonable throughout the course of each outbreak will vary. Examples of reasonable measures include:

- limiting visiting hours
- limiting the number of residents to whom the visitor has contact

- requiring anyone providing direct care (including visitors, other residents, etc) to wear the necessary PPE
- posting signs at entrances of facility and/or affected unit/area, discouraging visitors during the outbreak period; and
- notifying persons of the outbreak.

The above measures are seen to be reasonable and appropriate during the course of an outbreak. However, under outbreak conditions that present a greater risk to the resident population of the LTCH, more restrictive control measures may be required and occasionally there may be a conflict relating to public health unit recommended outbreak control measures. If the public health unit assesses the risk of not complying with outbreak control recommendations to be high, i.e. the probability of adverse health events to other residents, such as disease transmission, is high; the public health unit may have to consider a written order from the Medical Officer of Health to the licensee of the LTCH to ensure compliance with outbreak control measures. Under these circumstances, it is reasonable and necessary for the Medical Officer of Health to issue an order under the HPPA to the licensee of the LTCH, to:

- stop admissions to the LTCH
- restrict resident movement to and from the home, or
- bar visitors from the home

These are fairly significant measures, and presumably lesser measures would be discussed and implemented before admissions would be banned or visitors barred completely from the home.

Box 13. Updated Line listing: Resident and Staff Surveillance

Updated Line Listing: Resident and Staff Surveillance
<p>Resident Surveillance:</p> <ul style="list-style-type: none"> • New cases, with all appropriate information. • Names of residents who have recovered/recovery date. • The status of ill residents and any issues, such as worsening symptoms or complications. • Identification of the causative organism through lab results. • Transfers to/re-admission from acute-care hospitals • Deaths. <p>Staff Surveillance</p> <ul style="list-style-type: none"> • New staff cases, together with all appropriate information. • Names of staff who have recovered. • Expected return-to-work dates as determined in collaboration with the Public Health Unit.

5.1.1 Admissions and Returns from Absences

Generally, as an outbreak control measure, a public health unit would advise against admission of new residents to a facility or unit/floor experiencing an outbreak. Furthermore, public health units will need to be vigilant regarding admissions and those residents returning after absences to ensure that due diligence has been exercised by the LTCH in order to protect these residents and/or the residents with whom they may come into contact. Admissions and return from absence decisions should be made in consultation with the public health unit. See below for factors to consider.

A comprehensive approach to new admissions or return of residents from hospitals back to the LTCH requires consideration of a number of factors and careful judgment with respect to risk to individual residents as well as patients in the larger context of health care. Restricting admissions to a LTCH in outbreak may create a backlog in emergency departments or acute care, with a risk to patients in that system. On the other hand, admission of an unexposed resident into a LTCH may put them at risk and may lengthen the duration of the outbreak, with an impact to the larger resident population. A measured and considered approach is required in consultation with the Medical Officer of Health.

An applicant to a LTCH cannot be removed from a waitlist for a LTCH where an outbreak of disease prevents the applicant from moving into the home at the time that the CCAC offers to authorize the applicant's admission to the home.

New Admissions and Return of Non-cases

The admission of new residents and return of residents who have not been line-listed in the outbreak (i.e. are not known cases) is generally not advised. Changes to this control measure should be considered carefully with respect to resident safety and quality of life, as well as system capacity (See Appendix 15 – Transfer and Return Algorithm for use during Outbreaks). Members of the OMT from the LTCH and PHU should discuss the situation and carefully consider all relevant factors to assess if new admissions and/or return of non-cases are being considered, such as:

- What is the current status of the outbreak at the LTCH?
- Does the resident's attending physician at the hospital agree to the admission/return based on a review of the current health status of the resident? And are they aware of the outbreak?
- Is the resident protected from the outbreak pathogen through appropriate infection prevention and control measures? If the outbreak is due to influenza, is the resident protected by immunization and/or an antiviral drug?

- Are appropriate accommodations available for the returning resident? Will the resident return to an outbreak affected area of the LTCH?
- Has the resident or their substitute decision-maker been given information about the return to the LTCH?

5.1.2 Medical and Other Appointments

If possible, re-schedule all non-urgent medical and other appointments until after the outbreak is over.

5.1.3 Transfer to Hospital, Other Facility or Urgent Appointment

Notify the ICP at the receiving hospital/other facility regarding the details of the outbreak to ensure control measures are in place when the resident arrives and whether or not the transferring resident has been identified as a case or comes from an affected area.

It is **not** recommended to transfer residents from anywhere in the LTCH to another LTCH during an outbreak. The Outbreak Management Team or the Public Health Unit and the ICPs of both facilities may approve the transfer of residents on a case-by-case basis. The Provincial Transfer Authorization Centre, Ambulance, Community Care Access Centre or others must be notified about the outbreak when transfer is being arranged.

5.2 Control Measures for Staff and Volunteers

5.2.1 Well Staff

Minimize movement of staff, students and volunteers between affected/unaffected floors/units and consider cohorting – assigning some staff members to look after ill residents and others to look after well residents. Alternatively, consider assigning staff to a single unit/floor. Where possible, have recovering staff returning to work care for symptomatic residents. Staff, students or volunteers who also work at other health-care facilities, day-care centres and food premises should advise their employers that they have been working in a LTCH at which there is an outbreak. They should immediately stop working at all institutions/facilities if they develop symptoms of gastroenteritis illness. Depending on the policies of their employers, staff may be asked to not return to work until 48 hours after their last exposure at the outbreak institution. This period could be modified if the causative agent is known. Staff should change their uniforms between facilities and before leaving the affected facility.

5.2.2 ILL Staff

Staff and volunteers with gastrointestinal illness should not enter the LTCH, but should report the illness to their supervisor who will report it to the ICP, or follow the reporting procedure of the LTCH. Staff, students or volunteers with gastroenteritis should be excluded for at least 48 hours after symptoms have resolved. Once a specific causative agent is known, disease-specific exclusions apply. Discard all ready-to-eat foods (i.e. food not to be cooked) prepared by dietary staff that became ill while on shift.

5.3 Control Measures for Visitors and During Communal Activities

5.3.1 Modified Visitor Access

Complete closure of a LTCH to visitation is not permitted unless there is an order issued by the Medical Officer of Health as it may cause residents and visitors emotional hardship. Under exceptional circumstances, the Medical Officer of Health may assess the risk to be significant such that it requires complete closure to visitors. In these circumstances, an order from the MOH to the licensee is required to ensure compliance. It is important to note however that even under these circumstances, that there are exceptional personal circumstances under which barring visitors is neither ethical nor permitted. In these situations, the LTCH must ensure full compliance with infection control requirements. Furthermore, decisions to restrict visitors with or without an order of the MOH may be challenged and therefore need to be carefully considered and implemented. Visitation restrictions should be discussed by the OMT.

Post signs at all entrances, indicating the LTCH is experiencing an outbreak (Box 14).

Box 14. Notifying Visitors of Their Risk of Infection

Notifying Visitors of Their Risk of Infection*

Visitors should be advised that:

- There is a potential risk they could acquire illness while in the LTCH.
- They may pose a risk of introducing/reintroducing illness to the LTCH.

***Contact family members and advise them of their relative's illness.**

***If possible, keep a telephone list of frequent visitors and inform them of the outbreak.**

Well Visitors

Those who do visit during an outbreak must:

1. Perform hand hygiene when entering the LTCH, before entering and upon leaving the resident's room.
2. Visit residents only in their rooms and avoid communal areas.
3. Visit only one resident and leave the LTCH immediately after the visit, if both parents are in the home but in different locations, it is recommended that the healthy parent (non- outbreak case) be visited first.
4. Do not mingle with other residents.
5. Wear personal protective equipment as needed, especially if providing direct care, such as toileting, to the resident.

ILL Visitors

Advise visitors not to enter the LTCH if they have gastrointestinal symptoms (e.g. diarrhea, nausea or vomiting), respiratory symptoms (e.g. fever, cough or runny nose) or other communicable disease. This applies at all times, not only during outbreaks.

Visiting ILL Residents in Long-Term Care Homes

Place signs on the doors of ill residents' rooms or in other visible locations advising all visitors to check at the nursing station before entering the room. Visitors should visit ill residents only in their rooms. Discourage visitors from providing direct care to residents. If they choose to provide direct care, ensure they use appropriate PPE and perform careful and frequent hand hygiene without using the resident's sink. Visitors should be instructed about how to put on and remove PPE (Appendices 6, 7).

5.3.2 Communal Meetings and Other Activities

The following should be implemented during an outbreak:

- Reschedule communal meetings on the affected unit/floor. However, other meetings or activities may proceed in non-affected areas;
- Discontinue group outings from the affected unit/floor;
- The OMT should discuss restricting meetings or activities in the entire LTCH if the outbreak spreads to two or more units/floors;
- Do not permit visits by outside groups, such as entertainers, volunteer organizations and community groups;
- Conduct on-site programs such as physiotherapy and foot care for residents in their rooms, if possible. Proper precautions should be taken for ill residents; and
- Ensure there is no interaction between the affected floor/unit and participants in on-site child-care or other day programs.

6 Declaring the Outbreak Over and Termination of Outbreak Control Measures

The board of health (Public Health Unit) shall declare whether an outbreak is over in consultation with the LTCH. The board of health (Public Health Unit) shall use the most current available epidemiological data and best practices/guidance documents to determine when an outbreak can be declared over. The Medical Officer of Health retains the final authority to determine if an outbreak is over.

The outbreak may be declared over when the criteria in Section 6.1 are met. The OMT may identify ongoing surveillance needs after the outbreak has been declared over.

Ongoing surveillance needs include:

- Monitoring resident status, updating the line listing(s) and communicating with the Public Health Unit representative.
- Noting any deaths that occurred after the outbreak was declared over, including whether the deceased was part of the outbreak.
- Noting any further transmission among staff. Notify all those initially informed of the outbreak that the outbreak has ended.

6.1 Criteria for Declaring an Outbreak Over

The end of an outbreak is determined on a case-by-case basis. The specific period will be decided by the Public Health Unit in consultation with the LTCH and is based on the transmission risk. The specific period varies by micro-organism, but often is set at:

- No new cases after one infectious period plus one incubation period. For example, the most common type of enteric illness in LTCHs is Norovirus. As per Appendix 14 - Position Statement a Norovirus outbreak can be declared over after five days.
- 48 hours after the symptoms of the last case have resolved and all appropriate precautions were taken, there was no confirmed etiologic agent, norovirus was not suspected and Kaplan's Criteria were used.

In some circumstances, the Public Health Unit, in consultation with the LTCH, may decide that it is possible to resume some activities and discontinue some control measures during this period if:

- The last case was an isolated case on a unit. It may be possible to declare the outbreak over with the assurance that strict infection prevention and control measures will remain in place until all symptoms of the case have disappeared.
- The last case occurred in a staff person now excluded from the LTCH. It may be possible to declare the outbreak over once one incubation period has passed since the staff member was last present in the LTCH.

Since LTCHs have some sporadic gastrointestinal infection cases in non-outbreak situations, the OMT may need to attempt to differentiate between these sporadic cases and outbreak-associated cases in identifying the last outbreak-related resident case.

6.2 Review the Outbreak

Meet with the OMT to review management of the outbreak – what was handled well and what could be improved in managing future outbreaks. Recommendations should identify future preventive actions and/or necessary policy/protocol changes. They also should include possible reasons for the outbreak and steps to prevent similar outbreaks in the future. A representative from public health may attend this meeting, especially if there were concerns or issues.

6.3 Complete the Outbreak Investigation File

Review the outbreak file to ensure it contains full documentation including:

- Copies of laboratory and other results.
- Copies of all minutes and other communications.
- All other documents specific to the investigation/management of the outbreak, including notes/line lists.
- A summary report.

Infection prevention and control staff should store copies of all documents related to the outbreak. Public Health will also maintain file copies of all documents related to the outbreak and will report details of the outbreak to the MOHLTC's Public Health Division via an electronic database.

7 Investigation and Management of Food-Borne Outbreaks

Most enteric outbreaks in LTCHs are caused by viruses which are acquired through person-to-person transmission. However, other microbial agents can cause enteric disease in LTCHs as well. These include foodborne as well as waterborne agents such as Verotoxin-producing *Escherichia coli*, *Salmonella* species and *Listeria monocytogenes*. For this reason, it is important that the Public Health Units investigate the possibility that foodborne, waterborne or other agents are potential sources of any illnesses identified.

It is important to note that outbreaks spread from person-to-person still could have started with a point source such as ill kitchen staff or contaminated food or utensils. Investigation of a food-borne outbreak may be discontinued if, during early stages of an outbreak, a confirmed etiological agent is identified and that it is being transmitted person-to-person. The policies and procedures related to the investigation and management of food-borne outbreaks should be integrated into the LTCHs IPAC program.

7.1 Recommended Food Handling Policies and Procedures

Policies and procedures should be developed to cover all aspects of food handling. The Public Health Unit can offer guidance. Implementing appropriate policies and procedures can be instrumental in preventing outbreaks and controlling those that do occur.

Once policies and procedures are in place, consider scheduling regular in-service training for appropriate personnel and keep a record of the training – date, session name, presenter, training description and names of staff who attended.

Policies and procedures should include those related to food handling staff, records of food suppliers, retention of food samples, temperature records of potentially hazardous foods (PHF), catered food, food brought in by families, common kitchens, feeding assistance, dishwasher temperature/sanitizing records and kitchen equipment installation and maintenance.

7.1.1 Food-Handling Staff

Staff should be familiar with the exclusion criteria for food handlers outlined in the Infectious Diseases Protocols (2008). Public Health will provide recommendations on the screening of ill staff for enteric diseases if it is strongly suspected that the outbreak is food-borne. Depending upon the circumstances, screening of asymptomatic staff may also be considered.

7.1.2 Records of Food Suppliers

Food contaminated early in the food-production process could be widely distributed and thus become the source of many simultaneous outbreaks. Under the LTCHA, all LTCHs must maintain, and keep for one year, a record of purchases related to the food production system, including food delivery receipts. Public Health Units also require accurate records of food suppliers, including emergency contact information. Include suppliers of foods not generally considered potentially hazardous foods, such as fruits and vegetables, as these food items have also been involved in outbreaks.

7.1.3 Retaining Food Samples

Although not legislated, consideration should be given to implementing a policy of retaining 200 gram samples of ready-to-eat food items, potentially hazardous foods from each meal (Box 15).

7.1.4 Temperature Records of Potentially Hazardous Food

LTCHs should verify and record the final cooking, reheating and holding temperatures of potentially hazardous food. The LTCHs policies and procedures should clearly indicate which foods must be monitored and the documentation required.

Improper cooling procedures often cause food-borne illness. Public Health Units recommend documenting the cooling procedures used for potentially hazardous, prepared-in-advance food items that are cooled and reheated before being served. Food-preparation temperature records should be kept in accordance with the LTCHs retention policy, but not less than three months.

7.1.5 Catered Food and Food Brought in by Families

Often, food prepared off-site is available to residents. The LTCH should have clear policies outlining:

- Procedures to be followed if a resident wishes to have one or more meals catered (commercial caterers must be approved by the Public Health Unit).
- Advising direct care staff that food has been brought into the LTCH. Visitors should be advised not to offer food to other residents without direct care staff knowledge. Some residents are not able to consume certain foods due to underlying medical conditions.
- Which staff should be notified.
- The labeling requirements such as contents, resident's name and date prepared.
- Required storage procedures such as location, duration and holding temperature requirements.

Box 15. A Food Retention Policy

A Food Retention Policy
<p>Once a potential outbreak has been identified, food samples should not be discarded. What to include in your food retention policy:</p> <ul style="list-style-type: none">• Types of food to be retained.• Date of production.• Retention period (or date of discard).• Location of retained food samples.• Type of retention container.• Quantity of food to be retained.• Labeling requirements such as: date, type of food and time of meal. <p>Food samples should be kept frozen at or below – 18°C for 10 days.</p>

7.1.6 Common Servery or Kitchenette

Serveries or kitchenettes that are accessible to residents must meet the requirements of the Food Premises Regulation under the HPPA. These areas may pose special concerns since they can allow unrestricted access to food supplies that could lead to food contamination. The Food Premises Regulation is available at:

<https://www.ontario.ca/laws/regulation/170493>.

The LTCHs policies must be in compliance with the LTCHA, O. Reg 79/10 and the Food Premises Regulation. The policies must:

- Clearly define to which servery or kitchenette residents have access.
- Require all personally-owned resident food be labeled – contents, name of resident and date of preparation/decanting.
- State that anyone preparing or handling food, including cooking with residents as part of activities programming, must follow proper hand-hygiene procedures.

7.1.7 Feeding Assistance

The LTCH should have as part of their policy, a requirement for:

1. A person assisting residents with eating to perform hand hygiene before and after each meal.
2. A resident to be provided an opportunity to perform hand hygiene before and after each meal.
3. Staff from external agencies and volunteers not to enter the LTCH if they have symptoms of a communicable disease, particularly if they are experiencing diarrhea or vomiting.

7.1.8 Dishwasher Temperature and Sanitizing Records

Facilities must keep clear records of wash and rinse temperatures for each mechanical dishwasher. If the dishwasher has a low-temperature rinse and relies on chemical sanitizing, sanitizer concentration checks must be performed and documented daily, at minimum. Records must be dated, initialed and kept on site for at least three months. The LTCH and Public Health Unit staff should review the records during routine inspections and outbreak investigations.

7.1.9 Kitchen Equipment Installation and Maintenance

Keep records of any equipment that has been installed or repaired. Instructions on how each piece of equipment should be maintained, cleaned and sanitized should be readily available.

7.2 Food-borne Illness Investigation

The Public Health Unit is the lead in investigations to identify the source of food-borne outbreaks. These investigations rely heavily on the cooperation and assistance of LTCH staff. A public health inspector (PHI) or public health nurse (PHN) may take the following actions once an outbreak has been confirmed and food/water has not been ruled out as the source.

7.2.1 Questionnaires

The collection of information from residents, staff and others who may have been exposed to contaminated food or water is important in assessing the cause of an outbreak. Case-history information – especially symptoms, onset times and food consumed – will often identify the most likely causative food items and may help identify the most likely organisms.

7.2.2 Clinical Samples

In addition to collecting samples from ill residents, LTCH staff should be encouraged to submit stool samples if they experience symptoms included in the case definition. In certain circumstances, the Public Health Unit may request stool samples from asymptomatic (not ill) staff as well.

7.2.3 Food Service Operation

To complete the outbreak investigation, the PHI will require detailed information on:

- Foods eaten by residents, including foods with altered texture – such as pureed foods. How the food was prepared (menus, recipes and formulations), including records of cold- holding temperatures, final cooking temperatures and hot-holding temperatures, as well as the date and time each item was prepared.
- Purchase and inventory records.
- Processing records.
- Hazard Analysis Critical Control Point (HACCP) plans and records. Personnel responsible for each operation.
- List of suppliers.
- Records of personnel absenteeism due to illness.
- Equipment repair/maintenance records.
- Dishwashing and utensil washing records.
- Cleaning and sanitizing procedures and schedules.
- Resident meal seating plans.
- Records of bacteriological water sampling and water-supply maintenance if the LTCH is not on a municipal water supply.

7.2.4 HACCP Investigation

The PHI may decide that an assessment of food preparation using HACCP principles is needed. This may require the LTCH to prepare the suspect meal again. The goal of the HACCP audit is to identify improper food-handling practices, not to identify staff responsible for the outbreak. During the HACCP audit, the PHI may note:

- Temperature control, including storage, cooking, reheating and hot holding temperatures.
- Frequency and procedure on how staff wash their hands.
- The procedures used for wearing/removing gloves.
- Personal hygiene to prevent food contamination when they sneeze or cough.
- If staff understand the concept of cross-contamination and are knowledgeable on prevention methods.
- If staff are aware of the implications of improperly preparing potentially hazardous food.

7.2.5 Provision of Alternative Sources of Food and Water

Until the suspect food has been identified, the Public Health Unit may direct the LTCH to provide food and water from another source. If the LTCH provides food or meals for other organizations, such as Meals on Wheels, the Public Health Unit may order it to suspend those services until further notice.

7.2.6 Summary Report

A summary of the food safety investigation should be included in the outbreak investigation report. The summary should include detailed key inspection findings along with food and/or environmental sample results.

References and further information

- Baker GR et al. 2004. The Canadian adverse events study: the incidence of adverse events among hospital patients in Canada. *CMAJ* 170:1678-86.
- Berg DE et al. 1995. Control of nosocomial infections in an intensive care unit in Guatemala City. *Clin Infect Dis* 21:588-93.
- Boccia D et al. 2002. Waterborne outbreak of Norwalk-like virus gastroenteritis at a tourist resort, Italy. *Emerg Infect Dis* 8:563-568.
- Daly PB et al. 1992. Impact on knowledge and practice of a multiregional long-term care facility infection control training program. *Am J Infect Control* 20:225-33.
- Chadwick PR et al. 2000. Management of hospital outbreaks of gastroenteritis due to small round structured viruses. *Hosp Infect* 45:1-10.
- Community and Hospital Infection Control Association (CHICA). 1996. The role of the infection control practitioner—CHICA-Canada. *Can J Infect Control* 11:36-37.
- Gaynes R et al. 2001. Feeding back surveillance data to prevent hospital-acquired infections. *Emerg infect Dis* 7:295-298.
- Graves N et al. 2007. Economics and preventing hospital-acquired infection: broadening the perspective. *Infect Control Hosp Epidemiol* 28:178-84.
- Haley RW et al. 1985. The efficacy of infection surveillance and control programs in preventing nosocomial infections in U.S. hospitals. *Am J Epidemiol* 121:182-205.
- Harbarth S et al. 2003. The preventable proportion of nosocomial infections: an overview of published reports. *J Hosp Infect* 54:258-66.
- Kretzer EK and Larson EL. 1998. Behavioral interventions to improve infection control practices. *Am J Infect Control* 26:245-53.
- Lim J and Middleton D. 2003. Enteric outbreaks reported in Ontario, 2000-2002. *Public Health and Epidemiology Report: Ontario*. 14:202-207.
- Middleton D. 1999. Reported enteric outbreaks in Ontario 1999 *Public Health and Epidemiology Report: Ontario*. 10:137-143.
- Ministry of Health and Long-Term Care (MOHLTC). 1990. Enteric diseases screening recommendations and case management guidelines on foodhandlers and patient care workers. Currently being revised as *Guidelines for the Management of Enteric Diseases in Healthcare Workers, Food handlers and Day Care Staff and Attendees* (2010).
- MOHLTC. 2007a. Infection prevention and control core competency education for all health care providers. <http://govdocs.ourontario.ca/node/26660>

- MOHLTC. 2008. Just Clean Your Hands program. Released 2008.
<https://www.publichealthontario.ca/en/BrowseByTopic/InfectiousDiseases/JustCleanYourHands/Pages/Just-Clean-Your-Hands.aspx>
- MOHLTC. 2013. Roles and Responsibilities of Hospitals and Public Health Units for *Clostridium difficile* Infection (CDI) Reporting and Outbreak Management.
- Ontario Health Protection and Promotion Act, R.S.O. 1990, CHAPTER H.7. Includes R.R.O. 1990, Regulation 562 (Food Premises).
- Ontario Hospital Association and the Ontario Medical Association, Joint Committee on Communicable Disease Protocols. 2009. Enteric diseases surveillance protocol for Ontario hospitals. <https://www.oha.com/labour-relations-and-human-resources/health-and-safety/communicable-diseases-surveillance-protocols>
- Ontario Long-Term Care Homes Act, 2007, S.O. 2007, CHAPTER 8. Includes O. Reg. 79/10 (General).
- Ontario Occupational Health & Safety Act, R.S.O. 1990, c.0.1. Includes Health Care and Residential Facilities O. Reg. 67/93. 2007.
- Public Health Agency of Canada (PHAC). December 1997. Infection control guidelines: preventing the spread of Vancomycin-resistant Enterococci (VRE) in Canada. CCDR, Supplement, 23S8.
<http://www.phac-aspc.gc.ca/publicat/ccdr-rmtc/97vol23/23s8/vreindx.html>
- PHAC. 1998. Infection control guidelines: hand washing, cleaning, disinfection and sterilization in health care. CCDR, Supplement, 24S8.
<http://www.phac-aspc.gc.ca/publicat/ccdr-rmtc/98vol24/index.html>
- PHAC. July 1999. Infection control guidelines: routine practices and additional precautions for preventing the transmission of infection in health care. CCDR, Supplement, 25S4
<http://www.phac-aspc.gc.ca/publicat/ccdr-rmtc/99vol25/25s4/index.html>
- PIDAC. April 2014. Best practices for hand hygiene in all health care settings, 4th edition.
http://www.publichealthontario.ca/en/BrowseByTopic/InfectiousDiseases/PIDAC/Pages/PIDAC_Documents.aspx
- PIDAC. February 2013. (Annex A): Screening, Testing and Surveillance for Antibiotic Resistant Organisms (AROs) in all health care settings.
https://www.publichealthontario.ca/en/BrowseByTopic/InfectiousDiseases/PIDAC/Pages/Routine_Practices_Additional_Precautions.aspx
- PIDAC. July 2014. Best Practice for surveillance of health care-associated infections in patient and resident populations, 3rd edition.
https://www.publichealthontario.ca/en/BrowseByTopic/InfectiousDiseases/PIDAC/Pages/PIDAC_Documents.aspx

- PIDAC. May 2012. Best practices for environmental cleaning for prevention and control of infections in all health care settings, 2nd edition.
https://www.publichealthontario.ca/en/BrowseByTopic/InfectiousDiseases/PIDAC/Pages/PIDAC_Documents.aspx
- PIDAC. May 2013. Best practices for cleaning, disinfection and sterilization of medical equipment/devices in all health care settings, 3rd edition.
https://www.publichealthontario.ca/en/BrowseByTopic/InfectiousDiseases/PIDAC/Pages/Cleaning_Disinfection_Sterilization.aspx
- PIDAC. October 2012. Best practices for infection prevention and control programs in Ontario in all health care settings, 3rd edition.
https://www.publichealthontario.ca/en/BrowseByTopic/InfectiousDiseases/PIDAC/Pages/PIDAC_Documents.aspx
- PIDAC. November 2012. Routine practices and additional precautions in all health care settings, 3rd edition.
https://www.publichealthontario.ca/en/BrowseByTopic/InfectiousDiseases/PIDAC/Pages/Routine_Practices_Additional_Precautions.aspx
- Pittet D et al. 1999. Compliance with handwashing in a teaching hospital infection control program. *Ann Intern Med* 130:126-30.
- Pittet D et al. 2000. Effectiveness of a hospital-wide programme to improve compliance with hand hygiene. *Lancet* 356:1307-12.
- Rajda Z and Middleton D. 2004. Descriptive epidemiology of enteric outbreaks reported in Ontario, 2003. *Public Health and Epidemiology Report: Ontario*. 15:159-165.
- Rodriguez EM et al. 1996. Outbreak of viral gastroenteritis in a nursing home: importance of excluding ill employees. *Infect Control Hosp Epidemiol* 17:587-592.
- Siegel J et al. 2007. The healthcare infection control practices advisory committee. Guideline for isolation precautions: preventing transmission of infectious agents in healthcare settings. *Am J Infect Control* 35 [Suppl 2]:S64-164.
- Smith PW and Rusnak PG. 1997. Infection prevention and control in the long-term-care facility. *Am J Infect Control* 25:488-512.
- Stone PW et al. 2002. A systematic audit of economic evidence linking nosocomial infections and infection control interventions: 1990-2000. *Am J Infect Control* 30:145-52.
- Whitby M et al. 2007. Behavioural considerations for hand hygiene practices: the basic building blocks. *J Hosp Infect* 65:1-8.
- World Health Organization. 2005. WHO Guidelines on Hand Hygiene in Health Care (Advanced Draft): A Summary. World Alliance for Patient Safety, "Clean Hands are Safer Hands".
- Zingg W et al. 2005. Impact of an outbreak of norovirus infection on hospital resources. *Infect Control Hosp Epidemiol* 26:263-267.

Zoutman DE and Ford BD. 2005. The relationship between hospital infection surveillance and control activities and antibiotic-resistant pathogen rates. *Am J Infect Control* 33:1-5.

Zoutman DE et al. 2003. The state of infection surveillance and control in Canadian acute care hospitals. *Am J Infect Control* 31:266-72.

Appendix 1: Instructions for Feces Specimen Collection During Outbreaks

Bacterial, parasitic and viral agents may produce gastroenteritis. The Enteric Outbreak Kit has been designed for the investigation of these agents simultaneously at the beginning of an outbreak when the causative agent is unknown.

The use of this kit should be limited to the first 10 – 15 specimens collected from symptomatic persons at the onset of the outbreak. The Enteric Outbreak Kit includes three vials, each with a colour-coded cap:

- Green - Bacterial examination
- Yellow - Parasitology examination
- White - Viral and toxin examination

If the outbreak is suspected to be bacterial or viral in nature, do not collect/submit all three vials to the laboratory.

Make sure the laboratory requisition accurately reflects the examinations required. Each Enteric Outbreak Kit includes complete instructions on specimen collection, storage, and transportation. These instructions must be followed or the sample may not be tested.

For detailed information please refer to the Laboratory Guide for Gastroenteritis Outbreaks available at <http://www.publichealthontario.ca>

Image 1. Enteric Outbreak Kit



Image 2. Enteric Outbreak Kit (page 1)



Enteric Outbreak Kit

Instructions for the collection and transportation of clinical specimens for faeces cultures.

Obtain supplies, complete requisitions and label specimen vials

1. Remove the appropriate specimen collection vial(s) from the biohazard bag. Do not use expired kits.
2. Complete an "Enteric Disease Investigation Multiple Specimen Submission Form **OR** public health laboratory General Test Requisition". Include the outbreak number which is assigned by the local health unit.
3. On the main kit label located on the biohazard bag, fill in the required information with a ballpoint pen (press firmly). Peel this label off of the bag and place this label on the completed submission form in the area marked;
 - "Label" of the "Enteric Disease Investigation Multiple Specimen Submission Form".

OR

- If a public health laboratory General Test Requisition is used, fill in the required information with a ballpoint pen (press firmly).
4. Record the patient name on each of the vials used. Peel off one of the four corresponding kit numbered labels located on the biohazard bag. Place one label on each vial used.
5. Note: The specimen container is required to have the patient's full name and date of collection or two unique identifiers. The information on the specimen must be the same as the name and other identifier on the test requisition. Unmatched or mismatched specimens will not be processed.

Specimen collection

6. Faeces specimens that have been in contact with water in toilet are unacceptable.
 - a) Infants/Toddlers (not toilet trained) – Collect faeces sample (bowel movement) from soiled diaper or directly from "potty".
 - b) Older Children/Adults – Instruct the patient to defecate into a clean container.

↩ continued on reverse

F-SD-KI-006-001

Rev. May, 2010

Image 3. Enteric Outbreak Kit (page 2)



Place specimen in appropriate container

7. Using the spoon from each vial, select different sites of the faeces specimen, preferably blood, mucus or pus, and transfer to the vials as follows:
 - a) Bacteriology - GREEN-capped vial with red-coloured transport medium. A collecting device (*plastic spoon*) is fitted inside the cap. Add 2-3 spoonfuls of faeces. Mix into transport medium. Replace and tighten cap.
 - b) Parasitology – YELLOW-capped vial with clear liquid preservative and plastic spoon. Add faeces up to the line indicated. Mix well. Replace and tighten cap.
 - c) Virology/Toxin - WHITE-capped vial which is empty with a plastic spoon. Add faeces up to the line indicated. Replace and tighten cap.

Transportation

8. Place all vials in the biohazard bag. Place the completed test requisition in the outside pocket. Do not place the test requisition inside the biohazard bag containing the specimens.
9. Refrigerate specimens immediately. Do not freeze specimens.
10. Send specimens to the local Health Unit or laboratory as soon as possible.

STORAGE - Kits can be stored at room temperature until use. **DO NOT USE EXPIRED KITS.**

TO ORDER KITS or INFORMATION: Contact OAHPP order desk, Public Health Laboratory Toronto @ 416 235-5937 or order by fax @ 416 235-5753 or your local Public Health Laboratory.

Rev. May, 2010

Appendix 2: Routine Practices, Additional Precautions, and reporting requirements for selected pathogens or conditions associated with gastroenteritis outbreaks

(Adapted from PIDAC 2012). Contact = Contact Precautions; RP = Routine Practices; Infection Control refers to ICP or IPC program members.

Organism or Disease	Category	Type of	Single Room Needed	Duration of Precautions	Comments
Amoebiasis or Dysentery <i>Entamoeba histolytica</i>	Adult	RP	No		-Reportable Disease
	Incontinent or non-compliant adult	Contact	Yes		
Antibiotic Resistant Organisms (AROs) • not listed elsewhere		Contact may be indicated	May be indicated	Precautions, if required, are initiated and discontinued by Infection Control	-See also listings under ESBL and CRE. (Enterobacteriaceae)
Ascariasis or Roundworm infection <i>Ascaris lumbricoides</i>		RP	No		-No person-to-person transmission
Botulism	See Food Poisoning/Food-borne Illness				
Campylobacter species	Adult	RP	No		-Reportable Disease -Notify Infection Control
	Incontinent or non-compliant adult	Contact	Yes	Continue precautions until stools are formed	-Reportable Disease -Notify Infection Control

Organism or Disease	Category	Type of Precaution	Single Room Needed	Duration of Precautions	Comments
Cholera <i>Vibrio cholera</i>	Adult	RP	No		-Reportable Disease -Notify Infection Control
	Incontinent or non-compliant adult	Contact	Yes		-Reportable Disease -Notify Infection Control
Clostridium difficile		Contact	Yes	Continue precautions until formed stool for at least 2 consecutive days	-Outbreaks Reportable -Notify Infection Control. -Laboratory-confirmed cases may be cohorted
Coxsackievirus	See Enteroviral Infections				
Cryptosporidiosis	Adult	RP	No		-Reportable Disease -Notify Infection Control
	Incontinent or non-compliant adult	Contact	Yes		

Organism or Disease	Category	Type of Precaution	Single Room Needed	Duration of Precautions	Comments
Diarrhea	Acute infectious	See Gastroenteritis			
	Suspected <i>C. difficile</i> diarrhea	See <i>Clostridium difficile</i>			
Echinococcosis		RP	No		-No person-to-person transmission
Echovirus disease	See Enteroviral Infections				
Enterobiasis or Pinworm Disease <i>Enterobius vermicularis</i>		RP	No		-Transmission is faecal-oral directly or indirectly through contaminated articles eg., bedding.
Enteroviral Infections (Coxsackie viruses, Echo viruses)	Adult	RP	No		
Escherichia coli O157:H7	Incontinent or non-compliant adult	Contact	Yes	Continue precautions until stools are formed	-Reportable Disease -Notify Infection Control
	Adult	RP	No		

Organism or Disease	Category	Type of Precaution	Single Room Needed	Duration of Precautions	Comments
ENTEROBACTERIACEAE-RESISTANT Carbapenem-resistant Enterobacteriaceae (CRE)		Contact	Yes	Continue precautions for duration of hospitalization	-Notify Infection Control If readmitted, use Contact precautions
EXTENDED SPECTRUM BETA-LACTAMASE- PRODUCING ENTEROBACTERIACEAE (ESBL)		Contact may be indicated	May be indicated	Precautions, if indicated, are initiated and discontinued by Infection Control	-Notify Infection Control
Food poisoning or Food-borne Illness	Clostridium botulinum(botulism)	RP	No		-Reportable Disease -No person-to-person transmission
	Clostridium perfringens	RP	No		
	Salmonella or E. coli O157:H7 in incontinent or non-compliant adult if stool cannot be contained	Contact	Yes	Continue precautions until Salmonella and E. coli O157:H7 ruled out	-Reportable Disease -Notify Infection Control
	Other causes	RP	No		


Organism or Disease	Category	Type of Precaution	Single Room Needed	Duration of Precautions	Comments
Gastroenteritis	Acute infectious	Contact	Yes	Continue precautions until C.difficile and Norovirus or other viral agents ruled out	-Outbreaks are reportable -Notify Infection Control -See specific organism if identified
	Incontinent or non-compliant adult	Contact	Yes	Continue precautions for duration of illness	
Giardiasis <i>Giardia lamblia</i>	Adult	RP	No		-Reportable Disease
	Incontinent/non-compliant adult	Contact	Yes	Continue precautions until stools are formed	
Hand, Foot, & Mouth Disease	See Enteroviral Infections				
Histoplasmosis <i>Histoplasma capsulatum</i>		RP	No		-No person-to-person transmission
Hookworm Disease or Ancylostomiasis		RP	No		-No person-to-person transmission
Listeriosis <i>Listeria monocytogenes</i>		RP	No		-Reportable Disease

Organism or Disease	Category	Type of Precaution	Single Room Needed	Duration of Precautions	Comments
Norovirus		Contact	Yes	Continue precautions until 48 hours after resolution of symptoms	-Outbreaks Reportable -Notify Infection Control
Paratyphoid Fever <i>Salmonella paratyphi</i>		RP	No		-Reportable Disease
Pinworms	See Enterobiasis				
Rotavirus		Contact	Yes	Continue precautions until formed stool	
Roundworm Infection	See Ascariasis				
Salmonellosis <i>Salmonella species</i>	Adult	RP	No		-Reportable Disease -Notify Infection Control
	Incontinent/non-compliant adult	Contact	Yes	Continue precautions until formed stool	
Shigellosis <i>Shigella species</i>	See Gastroenteritis				
Strongyloidiasis <i>Strongyloides stercoralis</i>		RP	No		-May cause disseminated disease in immunocompromised

Organism or Disease	Category	Type of Precaution	Single Room Needed	Duration of Precautions	Comments
Tapeworm Infection <i>Diphyllobothrium latum</i> (fish), <i>Hymenolepis nana</i> and <i>Taenia saginata</i> (beef), <i>Taenia solium</i> (pork)		RP	No		-Autoinfection is possible
Typhoid Fever <i>Salmonella typhi</i>		RP	No		-Reportable Disease
Yersinia enterocolitica	See Gastroenteritis				

Appendix 3: Routine Practices for all Health Care Settings


Image 4. Routine Practices to be used with all patients

ROUTINE PRACTICES to be used with <u>ALL PATIENTS</u>	
	<p>Hand Hygiene</p> <p>Hand hygiene is performed using alcohol-based hand rub or soap and water:</p> <ul style="list-style-type: none"> ✓ Before and after each client/patient/resident contact ✓ Before performing invasive procedures ✓ Before preparing, handling, serving or eating food ✓ After care involving body fluids and before moving to another activity ✓ Before putting on and after taking off gloves and PPE ✓ After personal body functions (e.g., blowing one's nose) ✓ Whenever hands come into contact with secretions, excretions, blood and body fluids ✓ After contact with items in the client/patient/resident's environment
	<p>Mask and Eye Protection or Face Shield [based on risk assessment]</p> <ul style="list-style-type: none"> ✓ Protect eyes, nose and mouth during procedures and care activities likely to generate splashes or sprays of blood, body fluids, secretions or excretions. ✓ Wear within two metres of a coughing client/patient/resident.
	<p>Gown [based on risk assessment]</p> <ul style="list-style-type: none"> ✓ Wear a long-sleeved gown if contamination of skin or clothing is anticipated.
	<p>Gloves [based on risk assessment]</p> <ul style="list-style-type: none"> ✓ Wear gloves when there is a risk of hand contact with blood, body fluids, secretions, excretions, non-intact skin, mucous membranes or contaminated surfaces or objects. ✓ Wearing gloves is NOT a substitute for hand hygiene. ✓ Remove immediately after use and perform hand hygiene after removing gloves.
	<p>Environment and Equipment</p> <ul style="list-style-type: none"> ✓ All equipment that is being used by more than one client/patient/resident must be cleaned between clients/patients/residents. ✓ All high-touch surfaces in the client/patient/resident's room must be cleaned daily.
	<p>Linen and Waste</p> <ul style="list-style-type: none"> ✓ Handle soiled linen and waste carefully to prevent personal contamination and transfer to other clients/patients/residents.
	<p>Sharps Injury Prevention</p> <ul style="list-style-type: none"> ✓ NEVER RECAP USED NEEDLES. ✓ Place sharps in sharps containers. ✓ Prevent injuries from needles, scalpels and other sharp devices. ✓ Where possible, use safety-engineered medical devices.
	<p>Patient Placement/Accommodation</p> <ul style="list-style-type: none"> ✓ Use a single room for a client/patient/resident who contaminates the environment. ✓ Perform hand hygiene on leaving the room.

Appendix 4: Signage for entrance to room of a resident requiring Contact Precautions in non-acute care facilities

(PIDAC 2012)

Image 5. Contact Precautions - Non-acute Care Facilities

CONTACT PRECAUTIONS – Non-acute Care Facilities	
	<p>Hand Hygiene as per Routine Practices Hand hygiene is performed:</p> <ul style="list-style-type: none"> ✓ Before and after each resident contact ✓ Before performing invasive procedures ✓ Before preparing, handling, serving or eating food ✓ After care involving body fluids and before moving to another activity ✓ Before putting on and after taking off gloves and other PPE ✓ After personal body functions (e.g., blowing one's nose) ✓ Whenever hands come into contact with secretions, excretions, blood and body fluids ✓ After contact with items in the resident's environment ✓ Whenever there is doubt about the necessity for doing so ✓ Clean the resident's hands before he/she leaves his/her room
	<p>Resident Placement</p> <ul style="list-style-type: none"> ✓ Single room with own toileting facilities if resident hygiene is poor ✓ Door may remain open ✓ Perform hand hygiene on leaving the room or bed space
	<p>Gown [based on risk assessment]</p> <ul style="list-style-type: none"> ✓ Wear a long-sleeved gown for <u>direct care</u>* when skin or clothing may become contaminated
	<p>Gloves [based on risk assessment]</p> <ul style="list-style-type: none"> ✓ Wear gloves for <u>direct care</u>* ✓ Wearing gloves is NOT a substitute for hand hygiene ✓ Remove gloves on leaving the room or bed space and perform hand hygiene
	<p>Environment and Equipment</p> <ul style="list-style-type: none"> ✓ Dedicate routine equipment to the resident if possible (e.g., stethoscope, commode) ✓ Disinfect all equipment before it is used for another resident ✓ All high-touch surfaces in the resident's room must be cleaned at least daily
	<p>Visitors</p> <ul style="list-style-type: none"> ✓ Visitors must wear gloves and a long-sleeved gown if they will be in contact with other residents or will be providing <u>direct care</u>*, as required by Routine Practices ✓ Visitors must perform hand hygiene before entry and on leaving the room

* Direct Care: Providing hands-on care, such as bathing, washing, turning the resident, changing clothing, continence care, dressing changes, care of open wounds/lesions or toileting. Feeding and pushing a wheelchair are not classified as direct care.

Appendix 5: Hand Hygiene Fact Sheet for Health Care Settings

Image 6: Hand hygiene fact sheet for health care settings

Fact Sheet

Feuille de renseignements

PIDAC Provincial Infectious Diseases Advisory Committee

CCPMI Comité consultatif provincial des maladies infectieuses



Hand Hygiene Fact Sheet for Health Care Settings

In health care settings, hand hygiene is the single most important way to prevent infections.

Hand hygiene is the responsibility of all individuals involved in health care. Hand hygiene refers to removing or killing microorganisms on the hands as well as maintaining good skin integrity. There are two methods of removing/killing microorganisms on hands: washing with soap and running water or using an alcohol-based hand rub. Generally, the focus is on microorganisms that have been picked up by contact with patients/health care provider, contaminated equipment, or the environment (transient or contaminating bacteria).

Effective hand hygiene kills or removes microorganisms on the skin and maintains hand health.

ALCOHOL-BASED HAND RUB

Alcohol-based hand rub is the preferred method for decontaminating hands. Using alcohol-based hand rub is better than washing hands (even with an antibacterial soap) when hands are not visibly soiled.

However, hand washing with soap and running water must be performed when hands are visibly soiled. If running water is not available, use moistened towelettes to remove the visible soil, followed by alcohol-based hand rub.

HAND WASHING

Most transient bacteria present on the hands are removed during the mechanical action of washing, rinsing and drying hands. Hand washing with soap and running water must be performed when hands are visibly soiled.

WHEN SHOULD HAND HYGIENE BE PERFORMED?

Hand hygiene must be performed:

- Before and after contact with a patient
- Before performing invasive procedures
- Before preparing, handling, serving or eating food
- After care involving the body fluids of a patient (e.g. assisting patient to blow nose, toileting the patient or doing wound care) and before moving to another activity
- Before putting on and after taking off gloves
- After personal body functions, such as using the toilet or blowing one's nose
- Whenever a health care provider is in doubt about the necessity for doing so.
- When hands accidentally come into contact with secretions, excretions, blood and body fluids (hands must be washed with soap and running water)
- After contact with items in the patient's environment


FACTORS THAT INFLUENCE HAND HYGIENE

The following factors influence the effectiveness of hand hygiene:

- Condition of the skin— intact skin vs. presence of dermatitis, cracks, cuts or abrasions
- Nails: natural nails more than 3-4 mm (1/4-inch) long are difficult to clean, can pierce gloves and harbour more microorganisms than short nails
- Only nail polish in good condition is acceptable
- Artificial nails or nail enhancements are not to be worn by those giving patient care as they have been implicated in the transfer of microorganisms
- Jewellery - rings and bracelets hinder hand hygiene, and should not be worn for patient contact; rings increase the number of microorganisms present on hands and increase the risk of tears in gloves

Appendix 6: Recommended steps for putting on and taking off personal protective equipment (PPE)


(PIDAC 2012)



TAKING OFF PPE

1. Remove Gloves

- Remove gloves using a glove-to-glove/skin-to-skin technique
- Grasp outside edge near the wrist and peel away, rolling the glove inside-out
- Reach under the second glove and peel away
- Discard immediately into waste receptacle



2. Remove Gown

- Remove gown in a manner that prevents contamination of clothing or skin
- Starting at the neck ties, the outer, 'contaminated', side of the gown is pulled forward and turned inward, rolled off the arms into a bundle, then discarded immediately in a manner that minimizes air disturbance



6. Perform Hand Hygiene



3. Perform Hand Hygiene



5. Remove Mask/N95 Respirator

- Ties/ear loops/straps are considered 'clean' and may be touched with hands
- The front of the mask/respirator is considered to be contaminated
- Untie bottom tie then top tie, or grasp straps or ear loops
- Pull forward off the head, bending forward to allow mask/respirator to fall away from the face
- Discard immediately into waste receptacle



4. Remove Eye Protection

- Arms of goggles and headband of face shields are considered to be 'clean' and may be touched with the hands
- The front of goggles/face shield is considered to be contaminated
- Remove eye protection by handling ear loops, sides or back only
- Discard into waste receptacle or into appropriate container to be sent for reprocessing
- Personally-owned eyewear may be cleaned by the individual after each use



Appendix 7: Donning and removal of personal protective equipment (PPE)

Donning and Removal of Personal Protective Equipment

Personal protective equipment (PPE) is designed to protect health care providers in health care settings from exposure to potentially infectious material. When providing care to patients, these products protect the skin and mucous membranes of the eyes, nose, and mouth from exposure to blood, body and respiratory secretions.

Always perform hand hygiene immediately before donning and after removing PPE.

Always don your PPE before contact with patients.

Sequence for donning PPE

- perform hand hygiene
- gown (if applicable)
- mask
- eyewear
- gloves (if applicable)

1. How to don a gown

- opening is in the back
- fully cover torso from neck to knees, arms to end of wrists, and wrap around the back
- secure at neck and waist
- if gown is too small, use two gowns: the first ties in front, the second ties in back

2. How to don a mask

- secure on head with ear loops
- place over nose, mouth, and chin
- fit flexible nose piece over bridge
- adjust fit – snug to face and below chin

3. How to don eye protection

- position eyewear over eyes and secure to head using ear pieces

4. How to don gloves

- don gloves last
- insert hands into gloves
- extend gloves over gown cuffs (if wearing gown)

5. How to use gloved hands

- keep gloved hands away from face
- avoid touching or adjusting other PPE
- remove gloves if they become torn; perform hand hygiene before donning new gloves
- limit surfaces and items touched

Sequence for removing PPE

- all items must be removed and discarded carefully
- perform hand hygiene after gloves/gown removal before your hands go near your face (for removal of masks and eye protection) and after completion of PPE removal, and *any time you suspect your hands are contaminated during PPE removal.*

1. Glove removal

- outside of glove is 'dirty'; use glove-to-glove/skin-to-skin handling method
- grasp outside edge near wrist
- peel away from hand, turning glove inside out
- hold in opposite gloved hand
- slide ungloved finger under wrist of remaining glove
- peel off from inside, creating a bag for both gloves
- discard

2. Gown removal

- gown front and sleeves are 'dirty'; handle by inside/back of gown
- unfasten ties
- peel gown away from neck and shoulder
- turn contaminated outside surface toward the inside
- fold or roll into a bundle
- discard

3. Perform hand hygiene

4. Eyewear removal

- outside of eyepiece is 'dirty'; handle by earpieces
- grasp earpieces with ungloved hands
- pull away from face
- place in designated receptacle for reprocessing

5. Mask removal

- front of mask is 'dirty'; handle by ear-loops
- remove from face, in a downward direction, using ear-loops
- discard

6. Perform hand hygiene immediately after removing PPE.

For more information:

Visit our website at www.health.gov.on.ca/pandemic or call the Health Care Provider Hotline 1-866-212-2272. This phone line is open 24 hours a day, seven days a week.

Appendix 8: Cleaning checklist for an ill resident's room during an outbreak

Checklist for Daily Cleaning:

Use a fresh bucket, cloths and mop head. Always work from the cleanest areas to the dirtiest areas.

- Walls – check for visible soiling and clean if required
- Clean all horizontal surfaces and “touched” areas (tables, bed rails, call bells, work surfaces, mattresses/covers, doorknobs, sinks, light fixtures, chairs, phone, TV controls, soap dispensers)
- Clean bathroom, working from sink area to toilet area
- Clean floors

CHECKLIST FOR DISCHARGE CLEANING (TERMINAL CLEANING):

- Remove all dirty/used items (e.g. suction container, disposable items)
- Remove curtains before starting to clean the room
- Discard and replace the following:
 - Soap
 - Toilet paper
 - Alcohol-based hand rub
 - Glove box
 - Sharps container
- Use clean cloths, mop, supplies, and solution to clean the room
- Fill one bucket with the disinfectant so it is the correct strength
- Check to see if the mattress, pillows, and chairs are torn
- Report damaged items to your supervisor to have them replaced/repared
- DO NOT RE-USE CLOTHS: Use several cloths to clean a room. Use each cloth only once. Do not dip a cloth back into disinfectant solution after use and re-use it on another surface
- Always work from top to bottom and clean area to dirty
- Clean all surfaces and allow the disinfectant appropriate contact time with:
 1. Mattress
 2. Pillow
 3. BP cuff

4. Bedrails and bed controls
 5. Call bell
 6. Stethoscope and column
 7. Flow meters
 8. Suction tube and outer container
 9. Pull cord in washroom
 10. Over-bed table
 11. Inside drawers
 12. TV controls
 13. Soap dispenser
 14. Door handles
 15. Light switches
 16. Light cord
 17. Chair
 18. Phone
 19. Clean the following (and any other items that might be used on another resident) thoroughly before they are used by another resident
 20. Commodes/high toilet seat
 21. Wheelchairs
 22. Monitors
 23. IV poles
- Replace the sharps container when it is 2/3 full
 - Clean the outer canister of the suction container and red tubing
 - Remove all tape from the surfaces
 - Wash the sheepskin between residents
 - Wash the lift mesh or sheet between residents

Appendix 9: Sample screening-policies sign to use during outbreaks



The spread of germs that make you and others sick

Do not visit if you have a:

Cough Sneezing

Fever Cold

Diarrhea

Vomiting

Clean your hands with alcohol-based hand rub:

- when you arrive
- before leaving
- after coughing, sneezing or blowing nose

Appendix 11: Quick reference guide: suspected outbreak

This list has been compiled as a quick reference for your convenience. Some of these recommendations could require modifications due to particular situations. Please refer to the complete text for additional details. Implement any other measures deemed appropriate by your Infection Prevention and Control Professional or the Public Health Unit.

1. Notify Public Health

When two or more cases of suspect gastroenteritis occur within 48 hours. Remember the case definition for a Suspected Gastroenteritis Outbreak: two suspected cases of infectious gastroenteritis in a specific area, such as a home, unit or floor within 48 hours.

2. Implement Infection Prevention and Control Program

Isolation

- Confine ill residents to their rooms until at least 48 hours after symptoms have disappeared.
- Exclude any ill employees from work until at least 48 hours after symptoms have disappeared.
- Do not transfer well residents into a room with an ill resident.
- Do not admit new residents to the affected unit/floor.
- Do not re-admit residents who have not been line listed in the outbreak.
- You can re-admit residents who met the case definition.
- Suspend communal meetings and other activities in the outbreak area.
- Notify hospitals, other institutions, EMS and other persons or agencies listed in 4.2.5.
- Use cohort nursing or similar practices to reduce the potential of spread from ill to well residents.

Personal Protection

- Post signs advising visitors of the outbreak and the appropriate personal precautions they should take.
- Use gloves when caring for a resident who is incontinent of stool and has soiled themselves and their environment.
- Wear a gown if there is a strong possibility that clothing may become soiled.
- Use a mask and eye goggles or shield to protect the face from splashes if a resident has explosive diarrhea or projectile vomiting.
- Persons cleaning areas heavily contaminated with vomitus or feces should wear surgical masks.

- Review hand-hygiene procedures with all staff, including volunteers.
- Environment – for more information see Appendix 12.
- Increase the frequency of routine cleaning. Assume any and all frequently touched surfaces – door handles, railings, elevator buttons, tables, counters etc. – are contaminated.
- If possible, dedicate equipment to a particular symptomatic resident. If equipment must be shared, it must be cleaned and disinfected between residents.
- Clean then disinfect or discard equipment before it is taken from the resident's room.
- Clean and disinfect vomit and feces spillages promptly.
- If you are cleaning up vomit or diarrhea, you can reduce the risk of being infected by:
 - Wearing disposable gloves is recommended.
 - Using paper towels to soak up excess liquid, and putting the paper towels and any solid matter directly into a plastic garbage bag.
 - Cleaning the soiled area with soap and hot water. The same cleaning cloth or sponge should not be used to clean other areas as this may spread the virus.
 - Putting all cleaning cloths and disposable gloves into a plastic garbage bag.
 - Washing hands well using soap and warm water for a minimum of 15 seconds.
 - Reference: <https://www.healthlinkbc.ca/healthlinkbc-files/norovirus>
- Using care to avoid creating splashes or aerosols during clean up.
- Immediately covering spillage with dry disposable paper towels.
- Wearing appropriate personal protective equipment (gloves and gowns, and masks if splashing anticipated).
- Cleaning the area using disposable paper towels to remove all vomit and feces. Discarding used paper towels into plastic lined garbage.
- Removing all organic matter before disinfecting the area. Then disinfecting the area using a freshly prepared hypochlorite solution (1 part bleach to 9 parts water), or accelerated hydrogen peroxide or hospital grade disinfectant. Household cleaners other than bleach do not work for most of the viruses that cause vomiting and diarrhea.
- Allowing appropriate contact time. Wiping area dry. Removing personal protective equipment. Washing hands.
- Cleaning contaminated carpets and soft furnishings with hot water and detergent, or steam clean. Vacuum cleaning is not recommended.
- Terminal cleaning the resident environment when resident has been asymptomatic of gastroenteritis symptoms for 48 hours.
- Handling soiled laundry as little as possible, with minimum agitation, and transporting it in closed bags, prior to washing and drying.

Surveillance

- Maintain detailed surveillance and updated line lists (Appendices 11 and 14).
- Advise Public Health daily of new cases and other changes to the line list.
- Update case definition as appropriate.
- Collect stool specimens using outbreak kits provided by Public Health (Appendix 8).
- Advise Public Health when you have samples ready to ship.
- Refrigerate (4°C) specimens until shipped, unless otherwise advised by Public Health.

Appendix 12: Steps in the prevention and management of gastroenteritis outbreaks

Step	Component
1. Prevention Policies	<ul style="list-style-type: none"> • Policy Preparation • General • Food • Routine Practices and Additional Precautions • Infection Prevention and Control Committees • Education • Food Handling Practices
2. Preparation	<ul style="list-style-type: none"> • Outbreak Preparedness
3. Surveillance	<ul style="list-style-type: none"> • Surveillance
4. Identification	<ul style="list-style-type: none"> • Outbreak Detection • Infectious Gastroenteritis Case Definition • Outbreak Definition • Suspected Gastroenteritis Outbreak Definition • Gastroenteritis Outbreak Definition
5. Outbreak Management	<ol style="list-style-type: none"> 1. Assess the Outbreak 2. Implement General Outbreak Control Measures 3. Consult with the Public Health Unit 4. Declare an Outbreak 5. Notify Appropriate Individuals/Agencies 6. Hold an Initial Outbreak Management Team Meeting 7. Monitor the Outbreak on an Ongoing Basis 8. Declare the Outbreak Resolved 9. Complete the Outbreak Investigation File 10. Review the Outbreak

Appendix 13: Gastroenteritis outbreak: sample line-listing form

Image 7. Sample Line Listing Form

For Staff: Date Reported to Health Unit: _____ Outbreak: ____-____
 For Residents: Onset date of first case: _____

Institution Name: _____ Phone: _____
 Address: _____ Fax: _____
 Contact Person: _____

Causative Agent Isolated: _____ Phone: _____
 Public Health Investigator: _____ Fax: _____
 Case definition: _____

#	Last name	First name	Age	Sex	Occupation / Room	Diet	Onset date (d/m/y)	Symptoms*** (use legend below)					Specimen			Hospital * date (d/m/y)	Death** date (d/m/y)	Comments
													Date	Type	Results			

**hospitalized – admitted into a hospital due to outbreak, other hospitalizations should be recorded under comments.*
Symptoms:
 D = diarrhea F = fever
 N = nausea H = headache
 V = vomiting A = abdominal cramps
 Record name only once on the line listing form

Appendix 14: Position Statement: Recommended Length of Exclusion for Cases Associated with Norovirus Outbreaks and When to Declare Norovirus Outbreaks Over

Enteric, Zoonotic, and Vector-Borne Disease
Unit Public Health Division
Public Health Protection and Prevention
Branch Ministry of Health and Long-Term
Care September 2010

Acknowledgements

The Ministry of Health and Long-Term Care would like to acknowledge the contribution of the Expert Panel that developed Position Statement: Recommended Length of Exclusion for Cases Associated with Norovirus Outbreaks and When to Declare *Norovirus* Outbreaks Over.

Expert Panel

Dr. Allison McGeer

Microbiologist, Infectious Disease Consultant
Mt. Sinai Hospital

Ken Brown

Manager, Infectious Diseases Control Division
York Region Health Services

Amy Chiu

Senior Public Health Inspector
York Region Health Services

Dr. Irene Armstrong

Associate Medical Officer of Health
Toronto Public Health

Danielle Steinman

Manager, Communicable Disease Division
Peel Public Health

Debbie Valackis

Infection Control Specialist, Communicable Disease Division
Peel Public Health

Joseph Lam

Supervisor, Outbreak Management
Ottawa Public Health Anne-Luise Winter
Senior Epidemiologist, Surveillance and Epidemiology
Ontario Agency for Health Protection and Promotion

Dr. Dean Middleton

Senior Public Health Epidemiologist
Ontario Agency for Health Protection and Promotion

Yvonne Whitfield

Senior Program Consultant, Public Health Protection and Prevention Branch
Public Health Division, Ministry of Health and Long-Term Care

Dr. Erika Bontovics

Manager, Public Health Protection and Prevention Branch
Public Health Division, Ministry of Health and Long-Term Care

Post-Panel Contributors

Lisa Fortuna

Team Lead, Enteric, Zoonotic and Vector-Borne Diseases Unit
Public Health Division, Ministry of Health and Long-Term Care

Dr. Mark P. Nelder

Senior Program Consultant, Public Health Protection and Prevention Branch
Public Health Division, Ministry of Health and Long-Term Care

Background to issue

Over the past few years, the Ministry of Health and Long-Term Care (MOHLTC) has received numerous requests from public health units regarding extending the exclusion period for ill staff of high-risk settings suspected of having *Norovirus* from 48 to 72 hours after symptom resolution. Likewise, the MOHLTC has been requested to clarify the time required to declare outbreaks of *Norovirus* over.

While *Norovirus* is generally a mild and self-limited disease, deaths and other complications have occurred as a consequence of *Norovirus* infection. The greatest burden of illness occurs in the institutionalized elderly. This emphasizes the need for a consistent approach to *Norovirus* outbreak management.

Current Status

Currently, most jurisdictions rely on evidence in the scientific literature and working groups' decisions to inform their guidelines and best practice recommendations for *Norovirus*. The exclusion period currently used by many public health units in Ontario

is 48 hours. Criteria for declaring outbreaks over vary among Ontario jurisdictions with a range of 2 to 6 days after cessation of symptoms in the last resident case. These discussions and varying criteria highlight the need for a review of recommendations by the MOHLTC along with the development of a policy statement that can be applied consistently across Ontario.

Period of Exclusion

Summary of the Literature and Jurisdictional Scan

The majority of pre-1994 literature indicate Norovirus shedding occurs up to 48-72 hours after recovery (cessation of symptoms), including 24-48 hours,³ 48 hours,^{4,5} and 48-72 hours⁶. Based on this research, the standard exclusion period for cases in high-risk settings of Norovirus has been 48 hours after cessation of symptoms.^{7,8} High-risk settings include hospitals, nursing homes, food service establishments, and day nurseries.

Post-1994 literature indicates that Norovirus shedding occurs past the traditional 48-72 hour period, including 10 days⁹ and approximately 14 days, primarily because more sensitive laboratory testing methods became available.¹⁰ In a study authored by Robert L. Atmar et. al, using reverse transcription-PCR testing after experimental human infection, virus particles were detected in fecal samples 18 hours after inoculation and lasted a median of 4 weeks up to 8 weeks.¹¹ While viral shedding can occur for extended periods especially in immuno-compromised individuals, peak viral load in stool is highest in the first 25 to 72 hours post inoculation.¹² However the relationship between the amount of viral shedding and transmissibility has not been established. The relationship between shedding and transmissibility requires further study¹³.

The majority of jurisdictions examined employ a 48-hour exclusion policy. The policy is accepted across Canada including British Columbia and Newfoundland. Exclusion policies vary in the USA: 24 hours in California, 48 hours in Wisconsin, and 72 hours in Florida. Countries using a 48-hour exclusion policy include: Australia, Ireland, New Zealand, and the United Kingdom. While most jurisdictions use the 48-hour policy, very few base their policies on the scientific literature.

Recommendation: Period of Exclusion

Symptomatic staff associated with a Norovirus outbreak should be excluded for a minimum of 48 hours after symptom resolution.

Staff – Anyone conducting activities within an institution that will bring him/her into contact with residents. This includes all health care providers, support services such as housekeeping, food handlers, volunteers, and contract workers.

Declaring a Norovirus outbreak over

Summary of the Literature and Jurisdictional Scan

The literature regarding when to declare a Norovirus outbreak over is scant, with very

few recommendations based on scientific literature. Recommendations made by most jurisdictions are presumably based upon the advice of their respective expert and working groups. The World Health Organization recommends to "...declare the outbreak over when the number of new cases has returned to background levels."¹⁴

The decision of when to declare a Norovirus outbreak over varies greatly across jurisdictions. Public health units in Ontario use time periods of 2 to 6 days. In British Columbia, Fraser Health and the BC Provincial Infection Control Network declare a Norovirus outbreak over after 3 days have elapsed since the last resident case has resolved.

Recommendations: Declaring a Norovirus Outbreak Over

Traditionally, for other disease outbreaks, the outbreak is declared over when one incubation plus one period of communicability have passed with no new cases identified. Following this process, outbreaks of Norovirus can be declared over when there are no new cases after five days (one incubation period (2 days) plus one period of communicability (3 days)). Criteria for declaring a Norovirus outbreak over may be modified by the public health unit in collaboration with the Outbreak Management Team where applicable.

Note: It is the responsibility of the Medical Officer of Health or his/her designate to declare a Norovirus outbreak over. Declaring the outbreak over should be based on epidemiology of the outbreak in conjunction with conducting a risk assessment. Epidemiological evidence should indicate that the outbreak is under control, infection prevention and control measures have been appropriately applied and adhered to, and that the number of cases has diminished to baseline levels. Further, the risk assessment supports that decreasing some of the control measures will not prolong the outbreak nor compromise the well-being of the residents/patients, visitors or staff. For Norovirus outbreaks in long-term care homes, the decision to declare an outbreak over must be consistent with the recommendations outlined in A Guide to the Control of Gastroenteritis Outbreaks in Long-Term Care and Retirement Home Settings (MOHLTC 2010, pending publication). The occurrence of a single case with symptoms of gastroenteritis within the 5 day period should be investigated to determine if this is a single sporadic case, or linked with others.

References

Public Health Agency of Canada. 2006. Provincial/Territorial enteric outbreaks in Canada, 1996- 2003. June: J31-37.

Centers for Disease Control and Prevention. 2006. Norovirus in healthcare facilities: general information. Division of Healthcare Quality Promotion. December.

White KE, Osterholm MT, Mariotti JA, Korlath JA, Lawrence DH, Ristinen, Greenberg HB.

1986. A food borne outbreak of Norwalk virus gastroenteritis. Evidence for post-recovery transmission. *American Journal of Epidemiology* 124: 120-126.
- Reid JA, Caul EO, White DG, Palmer SR. 1988. Role of infected food handler in hotel outbreak of Norwalk-like viral gastroenteritis: implications for control. *Lancet* 2: 321-323.
- Patterson T, Hutchings P, Palmer S. 1993. Outbreak of SRSV gastroenteritis at an international conference traced to food handled by a post-symptomatic carrier. *Epidemiol. Infect.* 111: 157-162.
- Thornhill TS, Kalica AR, Wyatt RG, Kapikian AZ, Chanock RM. 1975. Pattern of shedding of the Norwalk particles in stools during experimentally induced gastroenteritis in volunteers as determined by immune electron microscopy. *Journal of Infectious Diseases* 132: 28-34.
- American Public Health Association. 2004. *Control of communicable disease*, 18th ed. Heyman D, editor. Washington, DC, USA.
- American Academy of Pediatrics. 2006. Caliciviruses. In: Pickering LK, Baker CJ, Long SS, McMillan JA, eds. *Red Book: 2006 Report of the Committee on Infectious Diseases*, 27th ed. Elk Grove Village, IL.
- Parashar UD, Dow L, Fankhauser RL, Humphrey CD, Miller J, Ando T, Williams KS, Eddy CR, Noel JS, Ingram T, Bresee JS, Monroe SS, Glass RI. 1998. An outbreak of viral gastroenteritis associated with consumption of sandwiches: implications for the control of transmission by food handlers. *Epidemiol. Infect.* 121: 615-621.
- Daniels NA, bergmire-Sweat DA, Schwab KJ, Hendricks KA, Reddy S, Rowe SM, Fankhauser RL, Monroe SS, Atmar RL, Glass RI, Mead P. 2000. A foodborne outbreak of gastroenteritis with Norwalk-like viruses: first molecular traceback to deli sandwiches contaminated during preparation. *Journal of Infectious Diseases* 181: 1467-1470.
- Atmar RL, Opekum AR, Gilger MA, Estes MK, Crawford SE, Neill FH, Graham DY. 2008.
- Norwalk virus shedding after experimental human infection. *Emerging Infectious Diseases* 14: 1553-1557.
- Graham DY, Jiang X, Tanaka T, Opekun AR, Madore HP, Estes MK. 1994. Norwalk virus infection of volunteers: new insights based on improved assays. *Journal of Infectious Diseases* 170: 34-43.
- Centers for Disease Control. 2010. *Guideline for the Prevention and Control of Norovirus Gastroenteritis Outbreaks in Healthcare Settings*. Division of Healthcare Quality Promotion. June 2010.
- World Health Organization. 2008. *Foodborne disease outbreaks: guidelines for investigation and control*. WHO Press, Geneva, Switzerland.

Appendix 15: Sample Transfer & Return Algorithm for Use During Outbreaks

Image 8. Sample Transfer and Return Algorithm During Outbreaks (1)

Transfers and Returns between Long-Term Care Homes and Hospitals during Outbreaks


The return of residents to a long-term care home (LTCH) during outbreaks is generally restricted in an effort to protect susceptible individuals from being exposed to respiratory infections such as influenza, and gastrointestinal infections such as norovirus. Returns to LTCHs are not automatically prohibited. They must be considered carefully with respect to resident safety and quality of life, as well as system capacity.

The sample algorithm provided here is a compilation of work done in southwestern, southeastern, and central eastern Ontario involving all relevant partner organizations. The tool is an outline of the process and factors to consider when making decisions about returning residents to their long term care homes after a hospital stay. It outlines opportunities for dialogue among the system partners who are involved in the care of residents: long-term care homes, hospitals, public health units, physicians, and of course, the residents themselves.

The sample algorithm provided here, may be used or adapted by stakeholders across Ontario who may not have documented their processes and considerations for transfers and returns between LTCHs and hospitals during an outbreak. It is intended to promote dialogue of key considerations. Users of this sample may modify it as appropriate to reflect their local practices, and should do so in consultation with relevant partners.

For more information, LTCHs can follow up with their Regional Infection Control Network and public health unit, or see the [Guide to the Control of Respiratory Infection Outbreaks in Long-Term Care Homes](#).

Image 9. Sample Transfer and Return Algorithm During Outbreaks (2)



Appendix 16: Epidemiology of selected pathogens associated with gastroenteritis outbreaks

AGENT	EPIDEMIOLOGY	INCUBATION PERIOD	SYMPTOMS
Bacterial			
Clostridium difficile	A leading cause of nosocomial diarrhea in hospitals and nursing homes	Variable	Watery diarrhea, fever, loss of appetite, nausea, abdominal pain/tenderness
Bacillus cereus	A well-recognized cause of food poisoning	1-6 hours where vomiting is predominant (toxin); 6-24 hours where diarrhea is predominant (infection)	An intoxication characterized in some cases by sudden onset of nausea and vomiting, in others by colic and diarrhea
Campylobacter spp. Verotoxigenic E.coli (O157:H7), plus other VT types Enteropathogenic E. coli Enterotoxigenic E. coli, Enteroinvasive E. coli Salmonella spp. Shigella spp. Vibrio cholerae and other spp. Yersinia enterocolytica and other spp.	Highest incidence is usually during spring and summer months	Varies with the organism; on average, 6 hours to a few days	Gastrointestinal upset, (bloody feces may be associated with Campylobacter spp. and certain enterotoxigenic E. coli including E. coli O157:H7)

AGENT	EPIDEMIOLOGY	INCUBATION PERIOD	SYMPTOMS
Clostridium botulinum	Human botulism is a serious but relatively rare intoxication caused by potent preformed toxins produced by Clostridium botulinum	Neurological symptoms usually appear within 12- 36 hours.	Early symptoms and signs are marked by fatigue, weakness and vertigo usually followed by blurred vision, dry mouth, and difficulty in swallowing and speaking
Clostridium perfringens	A well-recognized cause of food poisoning	≈ 10-12 hours	Intestinal disorder characterized by sudden onset of colic followed by diarrhea. Nausea is common. Vomiting and fever are usually absent
Staphylococcus aureus	Preformed toxin	≈ 2-4 hours	An intoxication of abrupt and sometimes violent onset, with severe nausea, cramps, vomiting, and prostration, often accompanied by diarrhea
Parasitic			
Cryptosporidium parvum	Worldwide occurrence	Not precisely known; thought to be 1-12 days with average of 7 days	Diarrhea, abdominal pain; more severe in immuno-compromised patients (duration up to 30 days)
Cyclosporacayetanensis	Worldwide occurrence; Ontario outbreaks food-related	≈ 7 days	Watery diarrhea, anorexia, weight loss, abdominal cramping, and flatulence (duration variable up to 4 weeks)

AGENT	EPIDEMIOLOGY	INCUBATION PERIOD	SYMPTOMS
Giardia lamblia	Worldwide occurrence; high prevalence in daycare centers where children are not toilet trained	3-5 days; median 7-10 days	Chronic diarrhea, abdominal cramping, bloating, and weight-loss (duration of 2-6 weeks)
Viral			
Astrovirus	Entire year. Mainly infects small children and the elderly	1-2 days	Diarrhea (duration 1-14 days)
Caliciviruses, including Norwalk-like viruses and Sapporo-like viruses	Winter, early spring. Most common cause of diarrhea in adults	1-2 days	Diarrhea, vomiting, and fever (duration 24-60 hours)
Enteric Adenovirus (types 40, 41)	Entire year Most common in young children	5-10 days	Watery Diarrhea, vomiting, fever (duration of 1-7 days)
Enterovirus	Late summer, fall. Most common in infants and young children	3-7 days	Diarrhea, rash, low grade fever (duration of up to 14 days)
Rotavirus	Late winter, early spring; infects primarily children; can cause nursing home outbreaks	2-3 days	Fever, vomiting, and diarrhea (duration of 3-9 days)
Other viruses including Torovirus and Picobirnaviruses	Not well studied. Children and adults including immunosuppressed patients	Not well established	Persistent and acute diarrhea in children

